

ACRP

BCRJ code:

0036

Cell Line:

ACRP

Species:

Homo sapiens

Vulgar Name: Human

Tissue:

Ovary

Morphology:

Epithelial

Disease:

Carcinoma

Growth Properties:

Adherent

Sex:

Female

Derivation:

The cell line was originated by intermittent treatment of cell line A2780 with cisplatin.

DNA Profile:

Amelogenin: X , X

D2S1338: 21,22

D19S433: 12

CSF1PO: 10,11

D13S317: 12,13

D16S539: 11,13

D5S818: 11,12

D7S820: 10

THO1: 6

TPOX: 8,10

vWA: 15,16

FGA: 19,23

D3S1358: 14,16

D18S51: 16,18

D8S1179: 15,17

D21S11: 28

Biosafety:

1

Additional info:

This cell is a ovary carcinoma cell line resistant to cisplatin, having a IC50 of 3uM to such drug.

Culture Medium:

RPMI-1640 medium modified to contain 2 mM L-glutamine, 10 mM HEPES, 1 mM sodium pyruvate, 4500 mg/L glucose, and 1500 mg/L sodium bicarbonate and fetal bovine serum to a final concentration of 10%.

Subculturing:

Remove medium, and rinse with PBS without calcium and magnesium.

Remove the solution and add an additional 1 to 2 mL of trypsin-EDTA solution. Allow the flask to sit at room temperature (or at 37°C) until the cells detach.

Add fresh culture medium, aspirate and dispense into new culture flasks.

NOTE: For more information on enzymatic dissociation and subculturing of cell lines consult Chapter 12 in Culture of Animal Cells, a manual of Basic Technique by R. Ian Freshney, 6th edition, published by Alan R. Liss, N.Y., 2010.

Medium Renewal: Every 2 to 3 days

Subcultivation ratio: 1:2 to 1:3

Culture Conditions:

Atmosphere: air, 95%; carbon dioxide (CO₂), 5%

Temperature: 37°C

Cryopreservation:

95% FBS + 5% DMSO (Dimethyl sulfoxide)

Thawing Frozen Cells:

SAFETY PRECAUTION: Is highly recommend that protective gloves and clothing always be used and a full face mask always be worn when handling frozen vials. It is important to note that some vials leak when submersed in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vessel exploding or blowing off its cap with dangerous force creating flying debris.

1. Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the Oring and cap out of the water. Thawing should be rapid (approximately 2 minutes).
2. Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
3. For cells that are sensitive to DMSO is recommended that the cryoprotective agent be removed immediately. Transfer the vial contents to a centrifuge tube containing 9.0 mL complete culture medium and spin at approximately 125 x g for 5 to 7 minutes.
4. Discard the supernatant and Resuspend cell pellet with the recommended complete medium (see the specific batch information for the culture recommended dilution ratio).
5. Incubate the culture in a appropriate atmosphere and temperature (see "Culture Conditions" for this cell line).

NOTE: It is important to avoid excessive alkalinity of the medium during recovery of the cells. It is suggested that, prior to the addition of the vial contents, the culture vessel containing the growth medium be placed into the incubator for at least 15 minutes to allow the medium to reach its normal pH (7.0 to 7.6).

References:

Sherman-Baust CA, Weeraratna AT, Rangel LB, Pizer ES, Cho KR, Schwartz DR, Shock T, Morin PJ. Remodeling of extracellular matrix through overexpression of collagen VI contributes to cisplatin resistance in ovarian cancer cells. *Cancer Cell*. 2003 Apr; 3 (4): 377-86.

Depositors:

LETICIA BATISTA AZEVEDO RANGEL; UFES