

Code:	0371
Cell Line:	TE-1
Species:	Homo sapiens Vulgar Name: Human
Tissue:	Esophagus
Morphology:	Epithelial-like
Disease:	Squamous Carcinoma
Growth Properties:	Adherent
Sex:	Male
Derivation:	Human cell line derived from esophageal cancer. Squamous cell carcinoma.
Biosafety:	1
Additional info:	
Culture Medium:	RPMI-1640 medium modified to contain 2 mM L-glutamine, 10 mM HEPES, 1 mM sodium pyruvate, 4500 mg/L glucose, and 1500 mg/L sodium bicarbonate with fetal bovine serum to a final concentration of 10%.

Subculturing:

Volumes used in this protocol are for 75 cm² flask; proportionally reduce or increase amount of dissociation medium for culture vessels of other sizes.

Remove and discard culture medium.

Briefly rinse the cell layer with PBS without calcium and magnesium to remove all traces of serum that contains trypsin inhibitor.

Add 1.0 to 3.0 mL of Trypsin-EDTA solution to flask and observe cells under an inverted microscope until the cell layer is dispersed (usually within 5 to 15 minutes).

Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal.

Add 6.0 to 8.0 mL of complete growth medium and aspirate cells by gently pipetting.

Transfer cell suspension to centrifuge tube and spin at approximately 125 x g for 5 to 10 minutes.

Discard supernatant and resuspend cells in fresh growth medium. Add appropriate aliquots of cell suspension to new culture vessels.

Place culture vessels in incubators at 37°C.

NOTE: For more information on enzymatic dissociation and subculturing of cell lines consult Chapter 12 in Culture of Animal Cells, a manual of Basic Technique by R. Ian Freshney, 6th edition, published by Alan R. Liss, N.Y., 2010.

Medium Renewal: twice per week

Subcultivation ratio: 1:8

Culture Conditions: Atmosphere: air, 95%; carbon dioxide (CO₂), 5% Temperature: 37°C

Cryopreservation: 95% FBS + 5% DMSO (Dimethyl sulfoxide)

Thawing Frozen Cells:

SAFETY PRECAUTION: It is highly recommended that protective gloves and clothing always be used and a full face mask always be worn when handling frozen vials. It is important to note that some vials leak when submerged in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vessel exploding or blowing off its cap with dangerous force creating flying debris.

1. Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the Oring and cap out of the water. Thawing should be rapid (approximately 2 minutes).
2. Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
3. For cells that are sensitive to DMSO it is recommended that the cryoprotective agent be removed immediately. Transfer the vial contents to a centrifuge tube containing 9.0 mL complete culture medium and spin at approximately 125 x g for 5 to 7 minutes.
4. Discard the supernatant and Resuspend cell pellet with the recommended complete medium (see the specific batch information for the culture recommended dilution ratio).
5. Incubate the culture in an appropriate atmosphere and temperature (see "Culture Conditions" for this cell line).

NOTE: It is important to avoid excessive alkalinity of the medium during recovery of the cells. It is suggested that, prior to the addition of the vial contents, the culture vessel containing the growth medium be placed into the incubator for at least 15 minutes to allow the medium to reach its normal pH (7.0 to 7.6).

References:

Nishihira T, Kasai M, Mori S, Watanabe T, Kuriya Y, Suda M, Kitamura M, Hirayama K, Akaishi T, Sasaki T Gann 1979; 70:575-84.

PubMed ID: 520749 Akaishi, T., Sekine, Y., Sanekata, K., Nishihira, T. and Kasai, M. "Characteristics of Growth and Effectiveness of Anti-Cancer Drugs on Tissue Cultured Human Esophageal Cancer Cells."Esophageal Cancer 1986; 35-38.Excerpta Medica, Current Clinical Practice Series 40. Kuriya Y, Kitamura M, Akaishi T, Hirayama K, Sekine Y, Nishihira T, Kasai M Tohoku J Exp Med 1983; 139:377-87.

PubMed ID: 6868074 Nishihira T, Hashimoto Y, Katayama M, Mori S, Kuroki T J Cancer Res Clin Oncol 1993; 119:441-9.

PubMed ID: 8509434 Hay, Park, Gazdar, Ed. Atlas of Human Tumor Cell Lines 1994; 269-282.Academic Press

Depositors:

Jolie Kiemlian Kwee - Instituto Nacional de Câncer