

## Data Sheet

PAGE 1/3

<b>BCRJ Code:</b>	0008
<b>Cell Line:</b>	25.9.17S
<b>Species:</b>	Mus musculus
<b>Vulgar Name:</b>	Mouse
<b>Tissue:</b>	Blood
<b>Cell Type:</b>	Hybridoma: B Lymphocyte
<b>Morphology:</b>	Lymphoblast
<b>Growth Properties:</b>	Suspension
<b>Derivation:</b>	This line was derived by fusing SP2/0-Ag14 myeloma cells with lymphoid cells from C3H mouse imunized with spleen cells from C3H.SW.
<b>Products:</b>	Immunoglobulin; monoclonal antibody; MHC antigens: ab against; I-Ab: ab against; IgG 2a
<b>Biosafety:</b>	1
<b>Additional Info:</b>	This hybridoma secretes a monoclonal antibody ( IgG2a kappa ) against MHC antigens, that reacts with I-Ab and I-Ad antigens. Cross reactions with H-2p and H-2q was also found. The specificity patterns does not correspond to any previously known I-a specificities.
<b>Culture Medium:</b>	RPMI-1640 medium modified to contain 2 mM L-glutamine, 1 mM sodium pyruvate, 4500 mg/L glucose and fetal bovine serum to a final concentration of 10%.
<b>Subculturing:</b>	Cultures can be maintained by addition of fresh medium. Alternatively, cultures can be established by centrifugation with subsequent resuspension at 1 x 10 <sup>5</sup> viable cells/mL. Maintain cultures at a cell concentration between 1 x 10 <sup>5</sup> and 1 x 10 <sup>6</sup> cells/mL. NOTE: Do not allow the cell concentration to exceed 1 x 10 <sup>6</sup> cells/mL.

## Data Sheet

PAGE 2/3

### Subculturing Medium Renewal:

Every 2 to 3 days

### Culture Conditions:

Atmosphere: air, 95%; carbon dioxide (CO<sub>2</sub>), 5% Temperature: 37°C

### Cryopreservation:

95% FBS + 5% DMSO (Dimethyl sulfoxide)

### Thawing Frozen Cells:

**SAFETY PRECAUTION:** It is strongly recommended to always wear protective gloves, clothing, and a full-face mask when handling frozen vials. Some vials may leak when submerged in liquid nitrogen, allowing nitrogen to slowly enter the vial. Upon thawing, the conversion of liquid nitrogen back to its gas phase may cause the vial to explode or eject its cap with significant force, creating flying debris.

1. Thaw the vial by gently agitating it in a 37°C water bath. To minimize contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 minutes).
2. Remove the vial from the water bath as soon as its contents are thawed and decontaminate it by dipping in or spraying with 70% ethanol. From this point, all operations must be performed under strict aseptic conditions.
3. For cells sensitive to DMSO, it is recommended to remove the cryoprotective agent immediately. Transfer the vial contents to a centrifuge tube containing 9.0 mL of complete culture medium and centrifuge at approximately 125 × g for 5 to 7 minutes.
4. Discard the supernatant and resuspend the cell pellet in the recommended complete medium (see specific batch information for the appropriate dilution ratio).
5. Incubate the culture under appropriate atmospheric and temperature conditions (see "Culture Conditions" for this cell line).

**NOTE:** It is important to avoid excessive alkalinity of the medium during cell recovery. To minimize this risk, it is recommended to place the culture vessel containing the growth medium in the incubator for at least 15 minutes before adding the vial contents. This allows the medium to stabilize at its normal pH (7.0 to 7.6).

### References:

J.Immunol. 126: 317-321, 1981.

### Depositors:

Alberto Nobrega, Universidade Federal do Rio de Janeiro

### Cellosaurus:

[CVCL D660](#)



