

Data Sheet

PAGE 1/3

BCRJ Code:	0023
Cell Line:	53-6.72
Species:	Rattus norvegicus (B cell); Mus musculus (myeloma), rat (B cell); mouse (myeloma)
Vulgar Name:	Mouse / Rat
Tissue:	Blood
Cell Type:	Hybridoma: B Lymphocyte
Morphology:	Lymphoblast
Growth Properties:	Suspension
Derivation:	Animals were immunized with mouse splenic or thymic lymphocytes. Spleen cells were fused with NS-1 myeloma cells.
Applications:	Can be used in a sandwich killing cytotoxicity assay when conjugated with arsanilic acid and used in conjunction with rabbit anti arsanilic acid antibody.
Products:	Immunoglobulin; monoclonal antibody; CD8 murine: ab against; IgM
Biosafety:	1
Culture Medium:	RPMI-1640 medium modified to contain 2 mM L-glutamine, 1 mM sodium pyruvate, 4500 mg/L glucose and fetal bovine serum to a final concentration of 10%.
Subculturing:	Cultures can be maintained by addition of fresh medium. Alternatively, cultures can be established by centrifugation with subsequent resuspension at 1×10^5 viable cells/mL. Maintain cultures at a cell concentration between 1×10^5 and 1×10^6 cells/mL. NOTE: Do not allow the cell concentration to exceed 1×10^6 cells/mL.

Data Sheet

PAGE 2/3

Subculturing Medium Renewal:

Every 2 to 3 days

Culture Conditions:

Atmosphere: air, 95%; carbon dioxide (CO₂), 5% Temperature: 37°C

Cryopreservation:

95% FBS + 5% DMSO (Dimethyl sulfoxide)

Thawing Frozen Cells:

SAFETY PRECAUTION: Is highly recommend that protective gloves and clothing always be used and a full face mask always be worn when handling frozen vials. It is important to note that some vials leak when submersed in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vessel exploding or blowing off its cap with dangerous force creating flying debris. 1. Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the Oring and cap out of the water. Thawing should be rapid (approximately 2 minutes). 2. Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions. 3. For cells that are sensitive to DMSO is recommended that the cryoprotective agent be removed immediately. Transfer the vial contents to a centrifuge tube containing 9.0 mL complete culture medium and spin at approximately 125 x g for 5 to 7 minutes. 4. Discard the supernatant and Resuspend cell pellet with the recommended complete medium (see the specific batch information for the culture recommended dilution ratio). 5. Incubate the culture in a appropriate atmosphere and temperature (see "Culture Conditions" for this cell line). **NOTE:** It is important to avoid excessive alkalinity of the medium during recovery of the cells. It is suggested that, prior to the addition of the vial contents, the culture vessel containing the growth medium be placed into the incubator for at least 15 minutes to allow the medium to reach its normal pH (7.0 to 7.6).

References:

Immunol. Rev. 47:63-90, 1979. Wilson ME, et al. Local suppression of IFN-gamma in hepatic granulomas correlates with tissue-specific replication of Leishmania chagasi. J. Immunol. 156: 2231-2239, 1996. PubMed: 8690913 Wong P, Rudensky AY. Phenotype and function of CD4+ T cells in mice lacking invariant chain. J. Immunol. 156: 2133-2142, 1996. PubMed: 8690902 Murray HW, et al. Multiple host defense defects in failure of C57BL/6 ep/ep (Pale Ear) mice to resolve visceral Leishmania donovani infection. Infect. Immun. 64: 161-166, 1996. PubMed: 8557335 Ledbetter JA, Herzenberg LA. Xenogeneic monoclonal antibodies to mouse lymphoid differentiation antigens. Immunol. Rev. 47: 63-90, 1979. PubMed: 398327

Depositors:

Banco de Células do Rio de Janeiro



fb.com/bancodecelulas/



[@bcrj_apobcam](https://www.instagram.com/bcrj_apobcam)



bcrj.org.br

FOR.PR.008.4-REV01-01-07-22

ATCC:

TIB-105