

Data Sheet

PAGE 1/3

BCRJ Code:	0356
Cell Line:	BV-2
Species:	Mus musculus
Vulgar Name:	Mouse; C57BL/6
Tissue:	Brain, microglial cells
Growth Properties:	Loosely attached and in suspension
Derivation:	This cell line was transformed by retroviruses
Biosafety:	2
Additional Info:	The BV-2 cells express the nuclear v-myc and the cytoplasmic v-raf oncogene products as well as the env gp70 antigen at the surface level; the BV-2 cells have morphological, phenotypical and functional markers of macrophages
Culture Medium:	RPMI-1640 medium modified to contain 2 mM L-glutamine, 4500 mg/L glucose with fetal bovine serum to a final concentration of 10%.
Subculturing:	Volumes used in this protocol are for 75 cm ² flask; proportionally reduce or increase amount of dissociation medium for culture vessels of other sizes. Remove culture medium with floating cells to a centrifuge tube. Briefly rinse the cell layer with PBS without calcium and magnesium to remove all traces of serum that contains trypsin inhibitor. Add 2.0 to 3.0 ml of Trypsin-EDTA solution to flask and observe cells under an inverted microscope until cell layer is dispersed (usually within 5 to 15 minutes). Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal. Add 6.0 to 8.0 ml of complete growth medium and aspirate cells by gently pipetting. Add appropriate aliquots of the cell suspension to new culture vessels. Incubate cultures at 37°C. NOTE: For more information on enzymatic dissociation and subculturing of cell lines consult Chapter 12 in Culture of Animal Cells, a manual of Basic Technique by R. Ian Freshney, 6th edition, published by Alan R. Liss, N.Y., 2010.

Data Sheet

PAGE 2/3

Subculturing Medium Renewal:

2 to 3 times per week

Subculturing Subcultivation Ratio:

1:5 to 1:10

Culture Conditions:

Atmosphere: air, 95%; carbon dioxide (CO₂), 5% Temperature: 37°C

Cryopreservation:

95% FBS + 5% DMSO (Dimethyl sulfoxide)

Thawing Frozen Cells:

SAFETY PRECAUTION: It is strongly recommended to always wear protective gloves, clothing, and a full-face mask when handling frozen vials. Some vials may leak when submerged in liquid nitrogen, allowing nitrogen to slowly enter the vial. Upon thawing, the conversion of liquid nitrogen back to its gas phase may cause the vial to explode or eject its cap with significant force, creating flying debris.

1. Thaw the vial by gently agitating it in a 37°C water bath. To minimize contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 minutes).
2. Remove the vial from the water bath as soon as its contents are thawed and decontaminate it by dipping in or spraying with 70% ethanol. From this point, all operations must be performed under strict aseptic conditions.
3. For cells sensitive to DMSO, it is recommended to remove the cryoprotective agent immediately. Transfer the vial contents to a centrifuge tube containing 9.0 mL of complete culture medium and centrifuge at approximately 125 × g for 5 to 7 minutes.
4. Discard the supernatant and resuspend the cell pellet in the recommended complete medium (see specific batch information for the appropriate dilution ratio).
5. Incubate the culture under appropriate atmospheric and temperature conditions (see "Culture Conditions" for this cell line).

NOTE: It is important to avoid excessive alkalinity of the medium during cell recovery. To minimize this risk, it is recommended to place the culture vessel containing the growth medium in the incubator for at least 15 minutes before adding the vial contents. This allows the medium to stabilize at its normal pH (7.0 to 7.6).

Data Sheet

PAGE 3/3

References:

Cell Immunol 1996;170:251-259 - PMID: 8660825 Infect Immunol
1992;60:3682-3688 - PMID: 1500177 J Neuroimmunol 1990;27:229-237 -
PMID: 2110186 J Neuroimmunol 1991;34:53-60 - PMID: 1894734 J
Neuroimmunol 1995;58:111-116 - PMID: 7730446 J Neuroimmunol
1998;93:102-107 - PMID: 10378873 J Neurosci Res 1992;31:616-621 - PMID:
1578513 Neuroimmunol 1991;32:249-257 - PMID: 2033118

Depositors:

José Antunes Rodrigues - Universidade de São Paulo

Cellosaurus:

[CVCL_0182](#)