

## Banco de Células do Rio de Janeiro

## Data Sheet

**PAGE 1/2** 

BCRJ Code:	0306
Cell Line:	C1498
Species:	Mus musculus
Vulgar Name:	Mouse
Cell Type:	Lymphoblast
Morphology:	Lymphoblast
Disease:	Acute Myeloid Leukemia
Growth Properties:	Suspension
Sex:	Female
Biosafety:	1
Addtional Info:	They are resistant to 0.1 mM cortisol. Antigen expression: The cells do not express B7-1.
Culture Medium:	Dulbecco's Modified Eagle's Medium (DMEM) modified to contain 2 mM L-glutamine, 4500 mg/L glucose and 10% of fetal bovine serum.
Subculturing:	Cultures can be maintained by addition of fresh medium or replacement of medium. Alternatively, cultures can beestablished by centrifugation with subsequent resuspension in fresh medium at 1 to 3 x 10e5 viable cells/mL. A saturation density of 1 to 2 X 10e6 is obtainable.
Subculturing Medium Renewal:	Every 2 to 3 days
Culture Conditions:	Atmosphere: air, 95%; carbon dioxide (CO2), 5% Temperature: 37°C
Cryopreservation:	95% FBS + 5% DMSO (Dimethyl sulfoxide)
f fb.com/bancodecelulas/ @ @l	pcrj_apabcam bcrj.org.br FOR.PR.008.4-REV01-01-07-22



## Banco de Células do Rio de Janeiro

## Data Sheet

**PAGE 2/2** 

Thawing Frozen Cells:	<ul> <li>SAFETY PRECAUTION: It is strongly recommended to always wear protective gloves, clothing, and a full-face mask when handling frozen vials. Some vials may leak when submerged in liquid nitrogen, allowing nitrogen to slowly enter the vial. Upon thawing, the conversion of liquid nitrogen back to its gas phase may cause the vial to explode or eject its cap with significant force, creating flying debris.</li> <li>1. Thaw the vial by gently agitating it in a 37°C water bath. To minimize contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 minutes).</li> <li>2. Remove the vial from the water bath as soon as its contents are thawed and decontaminate it by dipping in or spraying with 70% ethanol. From this point, all operations must be performed under strict aseptic conditions.</li> <li>3. For cells sensitive to DMSO, it is recommended to remove the cryoprotective agent immediately. Transfer the vial contents to a centrifuge tube containing 9.0 mL of complete culture medium and centrifuge at approximately 125 × g for 5 to 7 minutes.</li> <li>4. Discard the supernatant and resuspend the cell pellet in the recommended complete medium (see specific batch information for the appropriate dilution ratio).</li> <li>5. Incubate the culture under appropriate atmospheric and temperature conditions (see "Culture Conditions" for this cell line).</li> </ul> NOTE: It is important to avoid excessive alkalinity of the medium during cell recovery. To minimize this risk, it is recommended to place the culture vessel containing the growth medium in the incubator for at least 15 minutes before adding the vial contents. This allows the medium to stabilize at its normal pH (7.0 to 7.6).
References:	Ralph P. Retention of lymphocyte characteristics by myelomas and theta+-lymphomas: sensitivity to cortisol and phytohemagglutinin. J. Immunol. 110: 1470-1475, 1973.
Depositors:	EDGAR JULIAN PAREDES GAMERO/ALICE TEIXEIRA FERREIRA - UNIFESP
Cellosaurus:	<u>CVCL 3494</u>

f