

## Data Sheet

PAGE 1/3

<b>BCRJ Code:</b>	0291
<b>Cell Line:</b>	C3A [HepG2/C3A, derivative of Hep G2]
<b>Species:</b>	Homo sapiens
<b>Vulgar Name:</b>	Human
<b>Tissue:</b>	Liver
<b>Cell Type:</b>	Epithelial
<b>Morphology:</b>	Epithelial
<b>Disease:</b>	Hepatocellular Carcinoma
<b>Growth Properties:</b>	Adherent
<b>Sex:</b>	Male
<b>Age/Ethnicity:</b>	15 Year / Caucasian
<b>Derivation:</b>	C3A is clonal derivative of Hep G2 that was selected for strong contact inhibition of growth, high albumin production, high production of alpha fetoprotein (AFP) and ability to grow in glucose deficient medium.
<b>Applications:</b>	This cell line is a suitable transfection host.
<b>DNA Profile:</b>	Amelogenin: X,Y CSF1PO: 10,11 D13S317: 9,13 D16S539: 12,13 D5S818: 11,13 D7S820: 10 TH01: 9 TPOX: 8,9 vWA: 17
<b>Tumor Formation::</b>	No, in immunosuppressed mice Yes, in semisolid medium

## Data Sheet

PAGE 2/3

### Products:

alpha-fetoprotein (alpha fetoprotein); albumin; alpha2 macroglobulin (alpha-2-macroglobulin); alpha1 antitrypsin (alpha-1-antitrypsin); transferrin; alpha1 antichymotrypsin; (alpha-1-antichymotrypsin); haptoglobin; ceruloplasmin; plasminogen; complement (C4); C3 activator; fibrinogen; alpha1 acid glycoprotein (alpha-1 acid glycoprotein); alpha2 HS glycoprotein (alpha-2-HS-glycoprotein); beta lipoprotein (beta-lipoprotein); retinol binding protein (retinol-binding protein)

### Biosafety:

1

### Additional Info:

As the cells become confluent, there is a marked reduction in AFP secretion and an increase in albumin secretion. Gluconeogenesis activity is strongly oxygen dependent. The cells have nitrogen metabolizing activity comparable to perfused rat livers. There is no evidence of a Hepatitis B virus genome in this cell line.

### Culture Medium:

Dulbecco's Modified Eagle's Medium (DMEM) with 2 mM L-glutamine, 1.0 g/L glucose and 10% of fetal bovine serum.

### Subculturing:

Volumes used in this protocol are for 75 cm<sup>2</sup> flask; proportionally reduce or increase amount of dissociation medium for culture vessels of other sizes. Remove and discard culture medium. Briefly rinse the cell layer with PBS without calcium and magnesium to remove all traces of serum that contains trypsin inhibitor. Add 2.0 to 3.0 mL of Trypsin-EDTA solution to flask and observe cells under an inverted microscope until cell layer is dispersed (usually within 5 to 15 minutes). Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal. Add 6.0 to 8.0 mL of complete growth medium and aspirate cells by gently pipetting. Add appropriate aliquots of the cell suspension to new culture vessels. Incubate cultures at 37°C NOTE: For more information on enzymatic dissociation and subculturing of cell lines consult Chapter 12 in Culture of Animal Cells, a manual of Basic Technique by R. Ian Freshney, 6th edition, published by Alan R. Liss, N.Y., 2010.

### Subculturing Medium Renewal:

Twice per week

### Subculturing Subcultivation Ratio:

1:3 to 1:6 is recommended

### Culture Conditions:

Atmosphere: air, 95%; carbon dioxide (CO<sub>2</sub>), 5% Temperature: 37°C



## Data Sheet

PAGE 3/3

**Cryopreservation:** 95% FBS + 5% DMSO (Dimethyl sulfoxide)

**SAFETY PRECAUTION:** It is strongly recommended to always wear protective gloves, clothing, and a full-face mask when handling frozen vials. Some vials may leak when submerged in liquid nitrogen, allowing nitrogen to slowly enter the vial. Upon thawing, the conversion of liquid nitrogen back to its gas phase may cause the vial to explode or eject its cap with significant force, creating flying debris.

1. Thaw the vial by gently agitating it in a 37°C water bath. To minimize contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 minutes).
2. Remove the vial from the water bath as soon as its contents are thawed and decontaminate it by dipping in or spraying with 70% ethanol. From this point, all operations must be performed under strict aseptic conditions.
3. For cells sensitive to DMSO, it is recommended to remove the cryoprotective agent immediately. Transfer the vial contents to a centrifuge tube containing 9.0 mL of complete culture medium and centrifuge at approximately 125 × g for 5 to 7 minutes.
4. Discard the supernatant and resuspend the cell pellet in the recommended complete medium (see specific batch information for the appropriate dilution ratio).
5. Incubate the culture under appropriate atmospheric and temperature conditions (see "Culture Conditions" for this cell line).

**NOTE:** It is important to avoid excessive alkalinity of the medium during cell recovery. To minimize this risk, it is recommended to place the culture vessel containing the growth medium in the incubator for at least 15 minutes before adding the vial contents. This allows the medium to stabilize at its normal pH (7.0 to 7.6).

**References:** Kelly JH. Permanent human hepatocyte cell line and its use in a liver assist device (LAD). US Patent 5,290,684 dated Mar 1 1994

**Depositors:** Ana Cecília Oliveira - Fundação Oswaldo Cruz

