

Data Sheet

PAGE 1/3

BCRJ Code:	0063
Cell Line:	CCRF-CEM
Species:	Homo sapiens
Vulgar Name:	Human
Tissue:	Peripheral Blood
Cell Type:	T Lymphoblast
Morphology:	Lymphoblast
Disease:	Acute Lymphoblastic Leukemia
Growth Properties:	Suspension
Sex:	Female
Age/Ethnicity:	4 Year / Caucasian
Derivation:	Human lymphoblasts from peripheral blood of a child with acute leukemia.
Applications:	This cell line is a suitable transfection host.
DNA Profile:	Amelogenin: X CSF1PO: 10,11 D13S317: 11,12 D16S539: 10,13 D5S818: 12,13 D7S820: 9,13 THO1: 6,7 TPOX: 8 vWA: 17,19
Tumor Formation::	IN SYRIAN HAMSTER [PubMed: 4295047]
Products:	Genes Expressed: CD3; Homo sapiens, expressed ,CD5; Homo sapiens, expressed ,CD7; Homo sapiens, expressed ,CD4; Homo sapiens, expressed
Biosafety:	1

Data Sheet

PAGE 2/3

Culture Medium:

RPMI-1640 medium modified to contain 2 mM L-glutamine, 1 mM sodium pyruvate, 4500 mg/L glucose and fetal bovine serum to a final concentration of 10%.

Subculturing:

Cultures can be maintained by addition of fresh medium. Alternatively, cultures can be established by centrifugation with subsequent resuspension at 1×10^5 viable cells/mL. Maintain cultures at a cell concentration between 1×10^5 and 1×10^6 cells/mL. NOTE: Do not allow the cell concentration to exceed 1×10^6 cells/mL. Population Doubling Time about: 24-30 hours

Subculturing Medium Renewal:

Every 2 to 3 days

Culture Conditions:

Atmosphere: air, 95%; carbon dioxide (CO₂), 5% Temperature: 37°C

Cryopreservation:

95% FBS + 5% DMSO (Dimethyl sulfoxide)

Thawing Frozen Cells:

SAFETY PRECAUTION: It is highly recommended that protective gloves and clothing always be used and a full face mask always be worn when handling frozen vials. It is important to note that some vials leak when submerged in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vessel exploding or blowing off its cap with dangerous force creating flying debris. 1. Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the Oring and cap out of the water. Thawing should be rapid (approximately 2 minutes). 2. Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions. 3. For cells that are sensitive to DMSO is recommended that the cryoprotective agent be removed immediately. Transfer the vial contents to a centrifuge tube containing 9.0 mL complete culture medium and spin at approximately $125 \times g$ for 5 to 7 minutes. 4. Discard the supernatant and Resuspend cell pellet with the recommended complete medium (see the specific batch information for the culture recommended dilution ratio). 5. Incubate the culture in an appropriate atmosphere and temperature (see "Culture Conditions" for this cell line). NOTE: It is important to avoid excessive alkalinity of the medium during recovery of the cells. It is suggested that, prior to the addition of the vial contents, the culture vessel containing the growth medium be placed into the incubator for at least 15 minutes to allow the medium to reach its normal pH (7.0 to 7.6).

References:

Foley GE, et al. Continuous culture of human lymphoblasts from peripheral blood of a child with acute leukemia. *Cancer* 18: 522-529, 1965. PubMed: 14278051
Sandstrom PA, Buttke TM. Autocrine production of extracellular catalase prevents apoptosis of the human CEM T-cell line in serum-free medium. *Proc. Natl. Acad. Sci. USA* 90: 4708-4712, 1993. PubMed: 8506323
Adams RA. Formal discussion: the role of transplantation in the experimental investigation of human leukemia and lymphoma. *Cancer Res.* 27: 2479-2482, 1967. PubMed: 4170381
Uzman BG, et al. Morphologic variations in human leukemic lymphoblasts (CCRF-CEM cells) after long-term culture and exposure to chemotherapeutic agents. A study with the electron microscope. *Cancer* 19: 1725-1742, 1966. PubMed: 5224274
Adams RA, et al. Leukemia: serial transplantation of human leukemic lymphoblasts in the newborn Syrian hamster. *Cancer Res.* 27: 772-783, 1967. PubMed: 4295047
Miranda L, et al. Isolation of the human PC6 gene encoding the putative host protease for HIV-1 gp160 processing in CD4+ T lymphocytes. *Proc. Natl. Acad. Sci. USA* 93: 7695-7700, 1996. PubMed: 8755538
CCRF-CEM is a T lymphoblastoid cell line derived by G.E. Foley, et al. using cells obtained in November, 1964 from peripheral blood buffy coat of a 4-year-old Caucasian female with acute lymphoblastic leukemia.

Depositors:

Antonio Monteiro - Banco de Células do Rio de Janeiro

ATCC:

CCL-119

Cellosaurus:

[CVCL_0207](https://www.ebi.ac.uk/ces/entry/CVCL_0207)