

Data Sheet

| | |
|---------------------------|---|
| BCRJ Code: | 0071 |
| Cell Line: | COS-7 |
| Species: | Cercopithecus aethiops |
| Vulgar Name: | Monkey; African Green Monkey |
| Tissue: | Kidney |
| Cell Type: | Sv40 Transformed |
| Morphology: | Fibroblast |
| Growth Properties: | Adherent |
| Age/Ethnicity: | ADULT / |
| Applications: | This is an African green monkey kidney fibroblast-like cell line suitable for transfection by vectors requiring expression of SV40 T antigen. |
| Products: | T-antigen |
| Biosafety: | 2 |
| Culture Medium: | Dulbecco's Modified Eagle's Medium (DMEM) modified to contain 4 mM L-glutamine, 4500 mg/L glucose, 1 mM sodium pyruvate and fetal bovine serum to a final concentration of 10%. |

Subculturing:

Volumes used in this protocol are for 75 cm² flask; proportionally reduce or increase amount of dissociation medium for culture vessels of other sizes. T-75 flasks are recommended for subculturing this product. Remove and discard culture medium. Briefly rinse the cell layer with PBS without calcium and magnesium to remove all traces of serum that contains trypsin inhibitor. Add 2.0 to 3.0 mL of Trypsin-EDTA solution to flask and observe cells under an inverted microscope until cell layer is dispersed (usually within 5 to 15 minutes). Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal. Add 6.0 to 8.0 mL of complete growth medium and aspirate cells by gently pipetting. Add appropriate aliquots of the cell suspension to new culture vessels. Incubate cultures at 37°C. NOTE: For more information on enzymatic dissociation and subculturing of cell lines consult Chapter 12 in Culture of Animal Cells, a manual of Basic Technique by R. Ian Freshney, 6th edition, published by Alan R. Liss, N.Y., 2010.

**Subculturing
Medium Renewal:**

2 to 3 times per week

**Subculturing
Subcultivation
Ratio:**

1:4 to 1:8

Culture Conditions:

Atmosphere: air, 95%; carbon dioxide (CO₂), 5% Temperature: 37°C

Cryopreservation:

95% FBS + 5% DMSO (Dimethyl sulfoxide)

Thawing Frozen Cells:

SAFETY PRECAUTION: It is highly recommended that protective gloves and clothing always be used and a full face mask always be worn when handling frozen vials. It is important to note that some vials leak when submerged in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vessel exploding or blowing off its cap with dangerous force creating flying debris.

1. Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the Oring and cap out of the water. Thawing should be rapid (approximately 2 minutes).
2. Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
3. For cells that are sensitive to DMSO it is recommended that the cryoprotective agent be removed immediately. Transfer the vial contents to a centrifuge tube containing 9.0 mL complete culture medium and spin at approximately 125 x g for 5 to 7 minutes.
4. Discard the supernatant and Resuspend cell pellet with the recommended complete medium (see the specific batch information for the culture recommended dilution ratio).
5. Incubate the culture in an appropriate atmosphere and temperature (see "Culture Conditions" for this cell line).

NOTE: It is important to avoid excessive alkalinity of the medium during recovery of the cells. It is suggested that, prior to the addition of the vial contents, the culture vessel containing the growth medium be placed into the incubator for at least 15 minutes to allow the medium to reach its normal pH (7.0 to 7.6).

Gluzman Y. SV40-transformed simian cells support the replication of early SV40 mutants. *Cell* 23: 175-182, 1981. PubMed: 6260373 Fernandez LM, Puett D. Lys583 in the third extracellular loop of the lutropin/choriogonadotropin receptor is critical for signaling. *J. Biol. Chem.* 271: 925-930, 1996. PubMed: 8557706 Maestrini E, et al. A family of transmembrane proteins with homology to the MET-hepatocyte growth factor receptor. *Proc. Natl. Acad. Sci. USA* 93: 674-678, 1996. PubMed: 8570614 Campbell M, et al. The simian foamy virus type 1 transcriptional transactivator (Tas) binds and activates an enhancer element in the gag gene. *J. Virol.* 70: 6847-6855, 1996. PubMed: 8794326 Gonzalez Armas JC, et al. DNA immunization confers protection against murine cytomegalovirus infection. *J. Virol.* 70: 7921-7928, 1996. PubMed: 8892915 Jang SI, et al. Activator protein 1 activity is involved in the regulation of the cell type-specific expression from the proximal promoter of the human profilaggrin gene. *J. Biol. Chem.* 271: 24105-24114, 1996. PubMed: 8798649 Dittrich E, et al. A di-leucine motif and an upstream serine in the interleukin-6 (IL-6) signal transducer gp130 mediate ligand-induced endocytosis and down-regulation of the IL-6 receptor. *J. Biol. Chem.* 271: 5487-5494, 1996. PubMed: 8621406 Lee JH, et al. The proximal promoter of the human transglutaminase 3 gene. *J. Biol. Chem.* 271: 4561-4568, 1996. PubMed: 8626812 Chen Y, et al. Demonstration of binding of dengue virus envelope protein to target cells. *J. Virol.* 70: 8765-8772, 1996. PubMed: 8971005 Russell DW, Miller AD. Foamy virus vectors. *J. Virol.* 70: 217-222, 1996. PubMed: 8523528 Wright DA, et al. Association of human fas (CD95) with a ubiquitin-conjugating enzyme (UBC-FAP). *J. Biol. Chem.* 271: 31037-31043, 1996. PubMed: 8940097 Zhang J, et al. Dynamin and beta-arrestin reveal distinct mechanisms for G protein-coupled receptor internalization. *J. Biol. Chem.* 271: 18302-18305, 1996. PubMed: 8702465 Ozcelebi F, et al. Phosphorylation of cholecystokinin receptors expressed on chinese hamster ovary cells. *J. Biol. Chem.* 271: 3750-3755, 1996. PubMed: 8631990 Gibson S, et al. Functional LCK is required for optimal CD28-mediated activation of the TEC family tyrosine kinase EMT/ITK. *J. Biol. Chem.* 271: 7079-7083, 1996. PubMed: 8636141 Shaul PW, et al. Acylation targets endothelial nitric-oxide synthase to plasmalemmal caveolae. *J. Biol. Chem.* 271: 6518-6522, 1996. PubMed: 8626455 Ladner RD, et al. Identification of a consensus cyclin-dependent kinase phosphorylation site unique to the nuclear form of human deoxyuridine triphosphate nucleotidohydrolase. *J. Biol. Chem.* 271: 7752-7757, 1996. PubMed: 8631817 Wu X, et al. Demonstration of a physical interaction between microsomal triglyceride transfer protein and apolipoprotein B during the assembly of ApoB-containing lipoproteins. *J. Biol. Chem.* 271: 10277-10281, 1996. PubMed: 8626595 Hawes BE, et al. Phosphatidylinositol 3-kinase is an early intermediate in the G beta gamma-mediated mitogen-activated protein kinase signaling pathway. *J. Biol. Chem.* 271: 12133-12136, 1996. PubMed: 8647803 Arai H, Charo IF. Differential regulation of G-protein-mediated signaling by chemokine receptors. *J. Biol. Chem.* 271: 21814-21819, 1996. PubMed: 8702980 Hsieh CM, et al. APEG-1, a novel gene preferentially expressed in aortic smooth muscle cells, is down-regulated by vascular injury. *J. Biol. Chem.* 271: 17354-17359, 1996. PubMed: 8663449 Holtmann MH, et al. Multiple extracellular loop domains contribute critical determinants for agonist binding and activation of the secretin receptor. *J. Biol. Chem.* 271: 14944-14949, 1996. PubMed: 8663161

References:



Data Sheet

Depositors: Banco de Células do Rio de Janeiro

ATCC: CRL-1651

Cellosaurus: [CVCL_0224](#)