

Data Sheet

PAGE 1/4

BCRJ Code:	0100
Cell Line:	HeLa
Species:	Homo sapiens
Vulgar Name:	Human
Tissue:	Cervix
Cell Type:	Epithelial
Morphology:	Epithelial
Disease:	Adenocarcinoma
Growth Properties:	Adherent
Sex:	Female
Age/Ethnicity:	31 Year / Black
Applications:	These cells are a suitable transfection host. This cell line can be used to screen for Escherichia coli strains with invasive potential.
DNA Profile:	Amelogenin: X CSF1PO: 9,10 D13S317: 12,13,3 D16S539: 9,10 D5S818: 11,12 D7S820: 8,12 THO1: 7 TPOX: 8,12 vWA: 16,18
Virus Susceptibility::	Human adenovirus 3 Encephalomyocarditis virus Human poliovirus 1 Human poliovirus 2 Human poliovirus 3
Products:	Keratin
Biosafety:	2
Additional Info:	The cells are positive for keratin by immunoperoxidase staining. HeLa cells have been reported to contain human papilloma virus 18 (HPV-18) sequences. P53 expression was reported to be low, and normal levels of pRB (retinoblastoma suppressor) were found.
Culture Medium:	Dulbecco's Modified Eagle's Medium (DMEM) with 1% non-essential amino acids, 2 mM L-glutamine, 1 mM sodium pyruvate, 1.0 g/L glucose and 10% of bovine serum.
Subculturing:	Volumes used in this protocol are for 75 cm ² flask; proportionally reduce or increase amount of dissociation medium for culture vessels of other sizes. T-75 flasks are recommended for subculturing this product. Remove and discard culture medium. Briefly rinse the cell layer with PBS without calcium and magnesium to remove all traces of serum that contains trypsin inhibitor. Add 2.0 to 3.0 mL of Trypsin-EDTA solution to flask and observe cells under an inverted microscope until cell layer is dispersed (usually within 5 to 15 minutes). Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal. Add 6.0 to 8.0 mL of complete growth medium and aspirate cells by gently pipetting. Add appropriate aliquots of the cell suspension to new culture vessels. Incubate cultures at 37°C. NOTE: For more information on enzymatic dissociation and subculturing of cell lines consult Chapter 12 in Culture of Animal Cells, a manual of Basic Technique by R. Ian Freshney, 6th edition, published by Alan R. Liss, N.Y., 2010.
Subculturing Medium Renewal:	2 to 3 times per week
Subculturing Subcultivation Ratio:	1:2 to 1:6
Culture Conditions:	Atmosphere: air, 95%; carbon dioxide (CO ₂), 5% Temperature: 37°C
Cryopreservation:	95% FBS + 5% DMSO (Dimethyl sulfoxide)

Data Sheet

PAGE 2/4

SAFETY PRECAUTION: It is strongly recommended to always wear protective gloves, clothing, and a full-face mask when handling frozen vials. Some vials may leak when submerged in liquid nitrogen, allowing nitrogen to slowly enter the vial. Upon thawing, the conversion of liquid nitrogen back to its gas phase may cause the vial to explode or eject its cap with significant force, creating flying debris.

1. Thaw the vial by gently agitating it in a 37°C water bath. To minimize contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 minutes).

2. Remove the vial from the water bath as soon as its contents are thawed and decontaminate it by dipping in or spraying with 70% ethanol. From this point, all operations must be performed under strict aseptic conditions.

3. For cells sensitive to DMSO, it is recommended to remove the cryoprotective agent immediately. Transfer the vial contents to a centrifuge tube containing 9.0 mL of complete culture medium and centrifuge at approximately $125 \times g$ for 5 to 7 minutes.

4. Discard the supernatant and resuspend the cell pellet in the recommended complete medium (see specific batch information for the appropriate dilution ratio).

5. Incubate the culture under appropriate atmospheric and temperature conditions (see "Culture Conditions" for this cell line).

NOTE: It is important to avoid excessive alkalinity of the medium during cell recovery. To minimize this risk, it is recommended to place the culture vessel containing the growth medium in the incubator for at least 15 minutes before adding the vial contents. This allows the medium to stabilize at its normal pH (7.0 to 7.6).

Thawing Frozen Cells:



Data Sheet

PAGE 3/4

AOAC International Invasiveness by *Escherichia coli* of mammalian cells, microbiological method. Gaithersburg, MD:AOAC International;AOAC "Official Methods of Analysis of the AOAC International" 982.36. Baldi A, et al. Genomic structure of the human retinoblastoma-related Rb2/p130 gene. *Proc. Natl. Acad. Sci. USA* 93: 4629-4632, 1996. PubMed: 8643454 Gey GO, et al. Tissue culture studies of the proliferative capacity of cervical carcinoma and normal epithelium. *Cancer Res.* 12: 264-265, 1952. Chen TR. Re-evaluation of HeLa, HeLa S3, and HEp-2 karyotypes. *Cytogenet. Cell Genet.* 48: 19-24, 1988. PubMed: 3180844 Boshart M, et al. A new type of papillomavirus DNA, its presence in genital cancer biopsies and in cell lines derived from cervical cancer. *EMBO J.* 3: 1151-1157, 1984. PubMed: 6329740 Schneider-Gadicke A, Schwarz E. Different human cervical carcinoma cell lines show similar transcription patterns of human papillomavirus type 18 early genes. *EMBO J.* 5: 2285-2292, 1986. PubMed: 3023067 Schwarz E, et al. Structure and transcription of human papillomavirus sequences in cervical carcinoma cells. *Nature* 314: 111-114, 1985. PubMed: 2983228 Pater MM, Pater A. Human papillomavirus types 16 and 18 sequences in carcinoma cell lines of the cervix. *Virology* 145: 313-318, 1985. PubMed: 2992153 Yee C, et al. Presence and expression of human papillomavirus sequences in human cervical carcinoma cell lines. *Am. J. Pathol.* 119: 361-366, 1985. PubMed: 2990217 Scheffner M, et al. The state of the p53 and retinoblastoma genes in human cervical carcinoma cell lines. *Proc. Natl. Acad. Sci. USA* 88: 5523-5527, 1991. PubMed: 1648218 Jones HW Jr., et al. George Otto Gey. (1899-1970). The HeLa cell and a reappraisal of its origin. *Obstet. Gynecol.* 38: 945-949, 1971. PubMed: 4942173 Scherer WF, Hoogasian AF. Preservation at subzero temperatures of mouse fibroblasts (strain L) and human epithelial cells (strain HeLa). *Proc. Soc. Exp. Biol. Med.* 87: 480-487, 1954. PubMed: 13237281 Scherer WF, et al. Studies on the propagation in vitro of poliomyelitis viruses. IV. Viral multiplication in a stable strain of human malignant epithelial cells (strain HeLa) derived from an epidermoid carcinoma of the cervix. *J. Exp. Med.* 97: 695-710, 1953. PubMed: 13052828 Fang X, et al. Lysophosphatidylcholine stimulates activator protein 1 and the c-Jun N-terminal kinase activity. *J. Biol. Chem.* 272: 13683-13689, 1997. PubMed: 9153219 Bruder JT, Kovessdi I. Adenovirus infection stimulates the Raf/MAPK signaling pathway and induces interleukin-8 expression. *J. Virol.* 71: 398-404, 1997. PubMed: 8985363 Huber M, et al. Tyrosine phosphorylation events during coxsackievirus B3 replication. *J. Virol.* 71: 595-600, 1997. PubMed: 8985388 Olson JK, et al. Varicella-zoster virus Fc receptor gE glycoprotein: serine/threonine and tyrosine phosphorylation of monomeric and dimeric forms. *J. Virol.* 71: 110-119, 1997. PubMed: 8985329 Goodrum FD, Ornelles DA. The early region 1B 55-kilodalton oncoprotein of adenovirus relieves growth restrictions imposed on viral replication by the cell cycle. *J. Virol.* 71: 548-561, 1997. PubMed: 8985383 Löffler S, et al. CD9, a tetraspan transmembrane protein, renders cells susceptible to canine distemper virus. *J. Virol.* 71: 42-49, 1997. PubMed: 8985321 Hendricks DT, et al. FHIT gene expression in human ovarian, endometrial, and cervical cancer cell lines. *Cancer Res.* 57: 2112-2115, 1997. PubMed: 9187105 Hoppe HC, et al. Identification of phosphatidylinositol mannoside as a mycobacterial adhesin mediating both direct and opsonic binding to nonphagocytic mammalian cells. *Infect. Immun.* 65: 3896-3905, 1997. PubMed: 9284169 Rieder G, et al. Role of adherence in Interleukin-8 induction in *Helicobacter pylori*-associated gastritis. *Infect. Immun.* 65: 3622-3630, 1997. PubMed: 9284128 St. Geme JW, et al. Characterization of the genetic locus encoding *Haemophilus influenzae* type b surface fibrils. *J. Bacteriol.* 178: 6281-6287, 1996. PubMed: 8892830 Mansky LM. The mutation rate of human immunodeficiency virus type 1 is influenced by the vpr gene. *Virology* 222: 391-400, 1996. PubMed: 8806523 Döbelstein M, Shenk T. Protection against apoptosis by the vaccinia virus SPI-2 (B13R) gene product. *J. Virol.* 70: 6479-6485, 1996. PubMed: 8709286 Churchill MJ, et al. The rev-responsive element negatively regulates human immunodeficiency virus type 1 env mRNA expression in primate cells. *J. Virol.* 70: 5786-5790, 1996. PubMed: 8709194 Goodrum FD, et al. Adenovirus early region 4 34-kilodalton protein directs the nuclear localization of the early region 1B 55-kilodalton protein in primate cells. *J. Virol.* 70: 6323-6335, 1996. PubMed: 8709260 Kolanus W, et al. alphaLbeta2 integrin/LFA-1 binding to ICAM-1 induced by cytohesin-1 a cytoplasmic regulatory molecule. *Cell* 86: 233-242, 1996. PubMed: 8706128 Gan W, Rhoads RE. Internal initiation of translation directed by the 5'-untranslated region of the mRNA for eIF4G, a factor involved in the picornavirus-induced switch from cap-dependent to internal initiation. *J. Biol. Chem.* 271: 623-626, 1996. PubMed: 8557663 Koumenis C, Giaccia A. Transformed cells require continuous activity of RNA polymerase II to resist oncogene-induced apoptosis. *Mol. Cell. Biol.* 17: 7306-7316, 1997. PubMed: 9372962 You M, et al. ch-IAP1, a member of the inhibitor-of-apoptosis protein family, is a mediator of the antiapoptotic activity of the v-Rel oncoprotein. *Mol. Cell. Biol.* 17: 7328-7341, 1997. PubMed: 9372964 Hess MT, et al. Base pair conformation-dependent excision of benzo[a]pyrene diol epoxide-guanine adducts by human nucleotide excision repair enzymes. *Mol. Cell. Biol.* 17: 7069-7076, 1997. PubMed: 9372938 Bartz SR, et al. Human immunodeficiency virus type 1 cell cycle control: Vpr is cytostatic and mediates G2 accumulation by a mechanism which differs from DNA damage checkpoint control. *J. Virol.* 70: 2324-2331, 1996. PubMed: 8642659 Roller RJ, et al. Structure and function in the herpes simplex virus 1 RNA-binding protein US11: mapping of the domain required for ribosomal and nucleolar association and RNA binding in vitro. *J. Virol.* 70: 2842-2851, 1996. PubMed: 8627758 Jang SI, et al. Activator protein 1 activity is involved in the regulation of the cell type-specific expression from the proximal promoter of the human profilaggrin gene. *J. Biol. Chem.* 271: 24105-24114, 1996. PubMed: 8798649 Anderson SM, et al. Intercellular transfer of a glycosylphosphatidylinositol (GPI)-linked protein: release and uptake of CD4-GPI from recombinant adeno-associated virus-transduced HeLa cells. *Proc. Natl. Acad. Sci. USA* 93: 5894-5898, 1996. PubMed: 8650189 Ditttrich E, et al. A di-leucine motif and an upstream serine in the interleukin-6 (IL-6) signal transducer gp130 mediate ligand-induced endocytosis and down-regulation of the IL-6 receptor. *J. Biol. Chem.* 271: 5487-5494, 1996. PubMed: 8621406 Lee JH, et al. The proximal promoter of the human transglutaminase 3 gene. *J. Biol. Chem.* 271: 4561-4568, 1996. PubMed: 8626812 Duus KM, Grose C. Multiple regulatory effects of varicella-zoster virus (VZV) gL on trafficking patterns and fusogenic properties of VZV gH. *J. Virol.* 70: 8961-8971, 1996. PubMed: 8971025 Lieber A, et al. Recombinant adenoviruses with large deletions generated by cre-mediated excision exhibit different biological properties compared with first-generation vectors in vitro and in vivo. *J. Virol.* 70: 8944-8960, 1996. PubMed: 8971024 Yang C, Compans RW. Analysis of the cell fusion activities of chimeric simian immunodeficiency virus-murine leukemia virus envelope proteins: inhibitory effects of the R peptide. *J. Virol.* 70: 248-254, 1996. PubMed: 8523533 Schubert U, et al. The two biological activities of human immunodeficiency virus type 1 Vpu protein involve two separable structural domains. *J. Virol.* 70: 809-819, 1996. PubMed: 8551619 McKnight KL, Lemon SM. Capsid coding sequence is required for efficient replication of human rhinovirus 14 RNA. *J. Virol.* 70: 1941-1952, 1996. PubMed: 8627720 Lu FM, Lux SE. Constitutively active human notch 1 binds to the transcription factor CBF1 and stimulates transcription through a promoter containing a CBF1-responsive element. *Proc. Natl. Acad. Sci. USA* 93: 5663-5667, 1996. PubMed: 8643633 Yamaguchi Y, et al. Biochemical characterization and intracellular localization of the Menkes disease protein. *Proc. Natl. Acad. Sci. USA* 93: 14030-14035, 1996. PubMed: 8943055 Hewitt SM, et al. Differential function of Wilms' tumor gene WT1 splice isoforms in transcriptional regulation. *J. Biol. Chem.* 271: 8588-8592, 1996. PubMed: 8621487 Kotanides H, Reich NC. Interleukin-4-induced STAT6 recognizes and activates a target site in the promoter of the interleukin-4 receptor gene. *J. Biol. Chem.* 271: 25555-25561, 1996. PubMed: 8810328 Zhao Z, et al. Activation of mitogen-activated protein (MAP) kinase pathway by pervanadate, a potent inhibitor of tyrosine phosphatases. *J. Biol. Chem.* 271: 22251-22255, 1996. PubMed: 8703041 Quitschke WW, et al. The initiator element and proximal upstream sequences affect transcriptional activity and start site selection in the amyloid beta-protein precursor promoter. *J. Biol. Chem.* 271: 22231-22239, 1996. PubMed: 8703038 Hocking AM, et al. Eukaryotic expression of recombinant biglycan. *J. Biol. Chem.* 271: 19571-19577, 1996. PubMed: 8702651 Chen H, et al. Octamer binding factors and their coactivator can activate the murine PU.1 (spi-1) promoter. *J. Biol. Chem.* 271: 15743-15752, 1996. PubMed: 8663022 Macville M, et al. Comprehensive and definitive molecular cytogenetic characterization of HeLa cells by spectral karyotyping. *Cancer Res.* 59: 141-150, 1999. PubMed: 9892199 Dentistry--Preclinical evaluation of biocompatibility of medical devices used in dentistry--Test methods for dental materials. Geneva (Switzerland):International Organization for Standardization/ANSI;ISO ISO 7405:1997.

References:

Depositors: Departamento de Virologia, Instituto de Microbiologia, UFRJ, Rio de Janeiro.

Cellosaurus: [CVCL_0030](https://www.ebi.ac.uk/ces/entry/0030)

