

Data Sheet

PAGE 1/4

BCRJ Code:	0100
Cell Line:	HeLa
Species:	Homo sapiens
Vulgar Name:	Human
Tissue:	Cervix
Cell Type:	Epithelial
Morphology:	Epithelial
Disease:	Adenocarcinoma
Growth Properties:	Adherent
Sex:	Female
Age/Ethinicity:	31 Year / Black
Applications:	These cells are a suitable transfection host. This cell line can be used to screen for Escherichia coli strains with invasive potential.
DNA Profile:	Amelogenin: X CSF1PO: 9,10 D13S317: 12,13.3 D16S539: 9,10 D5S818: 11,12 D7S820: 8,12 THO1: 7 TPOX: 8,12 vWA: 16,18
Virus Succeptility::	Human adenovirus 3 Encephalomyocarditis virus Human poliovirus 1 Human poliovirus 2 Human poliovirus 3
Products:	Keratin
Biosafety:	2
Addtional Info:	The cells are positive for keratin by immunoperoxidase staining. HeLa cells have been reported to contain human papilloma virus 18 (HPV-18) sequences. P53 expression was reported to be low, and normal levels of pRB (retinoblastoma suppressor) were found.
Culture Medium:	Dulbecco's Modified Eagle's Medium (DMEM) with 1% non-essential amino acids, 2 mM L-glutamine, 1 mM sodium pyruvate, 1.0 g/L glucose and 10% of bovine serum.
Subculturing:	Volumes used in this protocol are for 75 cm2 flask; proportionally reduce or increase amount of dissociation medium for culture vessels of other sizes. T-75 flasks are recommended for subculturing this product. Remove and discard culture medium. Briefly rinse the cell layer with PBS without calcium and magnesium to remove all traces of serum that contains trypsin inhibitor. Add 2.0 to 3.0 mL of Trypsin-EDTA solution to flask and observe cells under an inverted microscope until cell layer is dispersed (usually within 5 to 15 minutes). Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal. Add 6.0 to 8.0 mL of complete growth medium and aspirate cells by gently pipetting. Add appropriate aliquots of the cell supension to new culture vessels. Incubate cultures at 37°C. NOTE: For more information on enzymatic dissociation and subculturing of cell lines consult Chapter 12 in Culture of Animal Cells, a manual of Basic Technique by R. Ian Freshney, 6th edition, published by Alan R. Liss, N.Y., 2010.
Subculturing Medium Renewal:	2 to 3 times per week
Subculturing Subcultivation Ratio:	1:2 to 1:6
Subculturing Subcultivation Ratio: Culture Conditions:	1:2 to 1:6 Atmosphere: air, 95%; carbon dioxide (CO2), 5% Temperature: 37°C

f

0



Data Sheet

PAGE 2/4

Thawing Frozen Cells:	 SAFETY PRECAUTION: It is strongly recommended to always wear protective gloves, clothing, and a full-face mask when handling frozen vials. Some vials may leak when submerged in liquid nitrogen, allowing nitrogen to slowly enter the vial. Upon thawing, the conversion of liquid nitrogen back to its gas phase may cause the vial to explode or eject its cap with significant force, creating flying debris. 1. Thaw the vial by gently agitating it in a 37°C water bath. To minimize contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 minutes). 2. Remove the vial from the water bath as soon as its contents are thawed and decontaminate it by dipping in or spraying with 70% ethanol. From this point, all operations must be performed under strict aseptic conditions. 3. For cells sensitive to DMSO, it is recommended to remove the cryoprotective agent immediately. Transfer the vial contents to a centrifuge tube containing 9.0 mL of complete culture medium and centrifuge at approximately 125 × g for 5 to 7 minutes. 4. Discard the supernatant and resuspend the cell pellet in the recommended complete medium (see specific batch information for the appropriate dilution ratio). 5. Incubate the culture under appropriate atmospheric and temperature conditions (see "Culture Conditions" for this cell line).
	NOTE: It is important to avoid excessive alkalinity of the medium during cell recovery. To minimize this risk, it is recommended to place the culture vessel containing the growth medium in the incubator for at least 15 minutes before adding the vial contents. This allows the medium to stabilize at its normal pH (7.0 to 7.6).

f



Data Sheet

PAGE 3/4

Standardization/ANSI;ISO ISO 7405:1997. Departamento de Virologia, Instituto de Microbiologia, UFRJ, Rio de Janeiro.	AOAC international Invasiones by Escherichia coli of mammalian esile, microbiologial method. Gathersburg, MD-AOAC international-AOAC -Official Methods of Analysis of the AOAC international-AOAC -Official Methods of Analysis of the AOAC international-AOAC -Official Methods of Analysis of the AOAC international-AOAC -Official Methods of ADA (et al. Genome Structure of the human retrinol standaro - related RAD/E3J Ogene. PC: Natl Acad. Sci. USA 93: 4023-4032, 1995. PubMet: 832446 - 4041. Tesus exolutions of the Les 33, and Hig-2 - Javartypse. CYtopenet: Cell Genet. 48: 10-2 - JBBS. PubMet: 310307 Edimarts 1, 2016. PubMet:
	biocompatibility of medical devices used in dentistryTest methods for dental materials. Geneva (Switzerland):International Organization for Standardization/ANSI;ISO ISO 7405:1997. Departamento de Virologia, Instituto de Microbiologia, UFRJ, Rio de Janeiro.

Cellosaurus:

f

Depositors:

References:

CVCL_0030



Data Sheet

PAGE 4/4

f

