

Data Sheet

BCRJ Code:	0273
Cell Line:	J774 1.6
Species:	Mus musculus
Vulgar Name:	Mouse
Tissue:	Reticulum
Cell Type:	Macrophage-Like
Morphology:	Macrophage
Disease:	Sarcoma
Growth Properties:	Adherent
Products:	nitric oxide
Biosafety:	1
Culture Medium:	Dulbecco's Modified Eagle's Medium (DMEM) with 1% non-essential amino acids, 2 mM L-glutamine, 1.0 g/L glucose and 10% of fetal bovine serum.
Subculturing:	Subcultures are prepared by scraping. For a 75 cm ² flask, remove all but 10 mL culture medium (adjust amount accordingly for other culture vessels). Dislodge cells from the flask substrate with a cell scraper; aspirate and add appropriate aliquots of the cell suspension into new culture vessels.
Subculturing Medium Renewal:	Twice per week
Subculturing Subcultivation Ratio:	1:6 to 1:8
Culture Conditions:	Atmosphere: air, 95%; carbon dioxide (CO ₂), 5% Temperature: 37°C

Data Sheet

PAGE 2/2

Cryopreservation: 95% FBS + 5% DMSO (Dimethyl sulfoxide)

SAFETY PRECAUTION: It is highly recommended that protective gloves and clothing always be used and a full face mask always be worn when handling frozen vials. It is important to note that some vials leak when submerged in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vessel exploding or blowing off its cap with dangerous force creating flying debris. 1. Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the Oring and cap out of the water. Thawing should be rapid (approximately 2 minutes). 2. Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions. 3. For cells that are sensitive to DMSO it is recommended that the cryoprotective agent be removed immediately. Transfer the vial contents to a centrifuge tube containing 9.0 mL complete culture medium and spin at approximately 125 x g for 5 to 7 minutes. 4. Discard the supernatant and Resuspend cell pellet with the recommended complete medium (see the specific batch information for the culture recommended dilution ratio). 5. Incubate the culture in an appropriate atmosphere and temperature (see "Culture Conditions" for this cell line). NOTE: It is important to avoid excessive alkalinity of the medium during recovery of the cells. It is suggested that, prior to the addition of the vial contents, the culture vessel containing the growth medium be placed into the incubator for at least 15 minutes to allow the medium to reach its normal pH (7.0 to 7.6).

Thawing Frozen Cells:

References: Guido Damiani et al. J. Exp. MEn. © The Rockefeller University Press. Volume 152:808-822, October 1980.

Depositors: Leonardo Nimrichter - Universidade Federal Do Rio De Janeiro

Cellosaurus: [CVCL HA24](#)

