

Data Sheet

PAGE 1/4

BCRJ Code:	0123
Cell Line:	JEG-3
Species:	Homo sapiens
Vulgar Name:	Human
Tissue:	Placenta
Morphology:	Epithelial
Disease:	Choriocarcinoma
Growth Properties:	Adherent
Applications:	This cell line is a suitable transfection host.
DNA Profile:	Amelogenin: X,Y CSF1PO: 11,12 D13S317: 9,11 D16S539: 13,14 D5S818: 10,11 D7S820: 10,12 THO1: 9,9.3 TPOX: 8 vWA: 16
Tumor Formation::	Yes, in nude mice; forms malignant tumor consistent with choriocarcinoma
Products:	Human chorionic gonadotrophin (hCG), somatomammotrophin
Biosafety:	1
Additional Info:	The cells are able to transform steroid precursors to estrone and estradiol.
Culture Medium:	Dulbecco's Modified Eagle's Medium (DMEM) with 1% non-essential amino acids, 2 mM L-glutamine, 1 mM sodium pyruvate, 1.0 g/L glucose with 10% of fetal bovine serum.

Data Sheet

PAGE 2/4

Remove medium, and rinse with PBS without calcium and magnesium. Remove the solution and add an additional 1 to 2 mL of trypsin-EDTA solution. Allow the flask to sit at room temperature (or at 37°C) until the cells detach. Add fresh culture medium, aspirate and dispense into new culture flasks. NOTE: For more information on enzymatic dissociation and subculturing of cell lines consult Chapter 12 in Culture of Animal Cells, a manual of Basic Technique by R. Ian Freshney, 6th edition, published by Alan R. Liss, N.Y., 2010.

Subculturing:**Subculturing Medium Renewal:**

2 to 3 days

Subculturing Subcultivation Ratio:

1:4 to 1:6

Culture Conditions:Atmosphere: air, 95%; carbon dioxide (CO₂), 5% Temperature: 37°C**Cryopreservation:**

95% FBS + 5% DMSO (Dimethyl sulfoxide)

SAFETY PRECAUTION: It is strongly recommended to always wear protective gloves, clothing, and a full-face mask when handling frozen vials. Some vials may leak when submerged in liquid nitrogen, allowing nitrogen to slowly enter the vial. Upon thawing, the conversion of liquid nitrogen back to its gas phase may cause the vial to explode or eject its cap with significant force, creating flying debris.

1. Thaw the vial by gently agitating it in a 37°C water bath. To minimize contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 minutes).
2. Remove the vial from the water bath as soon as its contents are thawed and decontaminate it by dipping in or spraying with 70% ethanol. From this point, all operations must be performed under strict aseptic conditions.
3. For cells sensitive to DMSO, it is recommended to remove the cryoprotective agent immediately. Transfer the vial contents to a centrifuge tube containing 9.0 mL of complete culture medium and centrifuge at approximately 125 × g for 5 to 7 minutes.
4. Discard the supernatant and resuspend the cell pellet in the recommended complete medium (see specific batch information for the appropriate dilution ratio).
5. Incubate the culture under appropriate atmospheric and temperature conditions (see "Culture Conditions" for this cell line).

NOTE: It is important to avoid excessive alkalinity of the medium during cell recovery. To minimize this risk, it is recommended to place the culture vessel containing the growth medium in the incubator for at least 15 minutes before adding the vial contents. This allows the medium to stabilize at its normal pH (7.0 to 7.6).

Fogh J, et al. Absence of HeLa cell contamination in 169 cell lines derived from human tumors. J. Natl. Cancer Inst. 58: 209-214, 1977. PubMed: 833871
Goodfellow M, et al. One hundred and twenty-seven cultured human tumor cell lines producing tumors in nude mice. J. Natl. Cancer Inst. 59: 221-226, 1977. PubMed: 77210034 . . Acta Endocrinol. Suppl. 153: 137-153, 1971.
Landers JE, et al. Translational enhancement of mdm2 oncogene expression in human tumor cells containing a stabilized wild-type p53 protein. Cancer Res. 57: 3562-3568, 1997. PubMed: 9270029
Roesler WJ, et al. The alpha-isoform of the CCAAT/enhancer-binding protein is required for mediating cAMP responsiveness of the phosphoenolpyruvate carboxykinase promoter in hepatoma cells. J. Biol. Chem. 271: 8068-8074, 1996. PubMed: 8626491
Kohler PO, Bridson WE. Isolation of hormone-producing clonal lines of human choriocarcinoma. J. Clin. Endocrinol. 32: 683-687, 1971.

Thawing Frozen Cells:

References:

Depositors:

ELAINE RODRIGUES MELLO; FACULDADE DE CIENCIAS FARMACEUTICAS - SP

Cellosaurus:

[CVCL_0363](#)