

## Data Sheet

PAGE 1/3

<b>BCRJ Code:</b>	0139
<b>Cell Line:</b>	L-132
<b>Species:</b>	Homo sapiens
<b>Vulgar Name:</b>	Human
<b>Tissue:</b>	Embryo
<b>Cell Type:</b>	Fibroblast
<b>Morphology:</b>	Fibroblast
<b>Disease:</b>	Normal
<b>Growth Properties:</b>	Adherent
<b>Derivation:</b>	This line was originally thought to be derived from embryonic lung tissue, but was subsequently found, based on isoenzyme analysis, HeLa marker chromosomes, and DNA fingerprinting, to have been established via HeLa cell contamination.
<b>DNA Profile:</b>	Amelogenin: X CSF1PO: 9,10 D13S317: 12,13.3 D16S539: 9,10 D5S818: 11,12 D7S820: 8,12 TH01: 7 TPOX: 8,12 vWA: 16,18
<b>Virus Susceptibility::</b>	hepatitis virus
<b>Products:</b>	Keratin
<b>Biosafety:</b>	2
<b>Culture Medium:</b>	Dulbecco's Modified Eagle's Medium (DMEM) with 1% non-essential amino acids, 2 mM L-glutamine, 1.0 g/L glucose and 10% fetal bovine serum.

## Data Sheet

PAGE 2/3

**Subculturing:**

Remove medium, and rinse with PBS without calcium and magnesium. Remove the solution and add an additional 1 to 2 mL of trypsin-EDTA solution. Allow the flask to sit at room temperature (or at 37°C) until the cells detach. Add fresh culture medium, aspirate and dispense into new culture flasks. NOTE: For more information on enzymatic dissociation and subculturing of cell lines consult Chapter 12 in Culture of Animal Cells, a manual of Basic Technique by R. Ian Freshney, 6th edition, published by Alan R. Liss, N.Y., 2010.

**Subculturing Medium  
Renewal:**

Twice per week

**Subculturing Subcultivation  
Ratio:**

1:2 to 1:8 is recommended

**Culture Conditions:**Atmosphere: air, 95%; carbon dioxide (CO<sub>2</sub>), 5% Temperature: 37°C**Cryopreservation:**

95% FBS + 5% DMSO (Dimethyl sulfoxide)

**SAFETY PRECAUTION:** It is strongly recommended to always wear protective gloves, clothing, and a full-face mask when handling frozen vials. Some vials may leak when submerged in liquid nitrogen, allowing nitrogen to slowly enter the vial. Upon thawing, the conversion of liquid nitrogen back to its gas phase may cause the vial to explode or eject its cap with significant force, creating flying debris.

1. Thaw the vial by gently agitating it in a 37°C water bath. To minimize contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 minutes).
2. Remove the vial from the water bath as soon as its contents are thawed and decontaminate it by dipping in or spraying with 70% ethanol. From this point, all operations must be performed under strict aseptic conditions.
3. For cells sensitive to DMSO, it is recommended to remove the cryoprotective agent immediately. Transfer the vial contents to a centrifuge tube containing 9.0 mL of complete culture medium and centrifuge at approximately  $125 \times g$  for 5 to 7 minutes.
4. Discard the supernatant and resuspend the cell pellet in the recommended complete medium (see specific batch information for the appropriate dilution ratio).
5. Incubate the culture under appropriate atmospheric and temperature conditions (see "Culture Conditions" for this cell line).

**NOTE:** It is important to avoid excessive alkalinity of the medium during cell recovery. To minimize this risk, it is recommended to place the culture vessel containing the growth medium in the incubator for at least 15 minutes before adding the vial contents. This allows the medium to stabilize at its normal pH (7.0 to 7.6).

Didier ES, et al. Characterization of Encephalitozoon (Septata) intestinalis isolates cultured from nasal mucosa and bronchoalveolar lavage fluids of two AIDS patients. J. Eukaryot. Microbiol. 43: 34-43, 1996. PubMed: 8563708  
Erfle V, Mellert W. Human cell line LC5 and its use. US Patent 5,582,967 dated Dec 10 1996 . . Fed. Proc. 19: 386, 1960. . . Science 133: 2059, 1961.  
Tresnan DB, et al. Feline aminopeptidase N serves as a receptor for feline, canine, porcine, and human coronaviruses in serogroup I. J. Virol. 70: 8669-8674, 1996. PubMed: 8970993

## Thawing Frozen Cells:

## References:

## Depositors:

Jose Paulo Leite, Instituto Oswaldo Cruz, Rio de Janeiro.

## Cellosaurus:

[CVCL\\_1908](#)