

Data Sheet

BCRJ Code: 0315

Cell Line: MAC-T

Species: Bos taurus

Vulgar Name: Cow, Bovine

Tissue: Udder

Cell Type: Epithelial

Morphology: Epithelial

Growth Properties: Adherent

Applications:

Development of the MAC-T bovine mammary epithelial cell line by stable transfection with simian virus-40 large T-antigen should greatly assist study of possible intrinsic (local) and extrinsic (systemic) factors regulating bovine mammary epithelial cell development, differentiation, and function.

Biosafety:

2

Additional Info:

MAC-T cells were established from primary bovine mammary alveolar cells by stable transfection with SV-40 large T-antigen. MAC-T cells show a population doubling time of approximately 17 h and have been cultured more than 350 passages without showing any sign of senescence. They show the characteristic "cobblestone" morphology of epithelial cells when grown on plastic substratum. Differentiation was induced by augmenting cell-cell interaction on a floating collagen gel in the presence of prolactin.

Culture Medium:

Dulbecco's Modified Eagle's Medium (DMEM) modified to contain 2 mM L-glutamine, 4500 mg/L glucose, 5 ug/mL insulin, 1 ug/mL hydrocortisone and 10% of fetal bovine serum.

Subculturing:

Remove and discard culture medium. Briefly rinse the cell layer with PBS without calcium and magnesium to remove all traces of serum that contains trypsin inhibitor. Add 2.0 to 3.0 mL of Trypsin-EDTA solution to flask and observe cells under an inverted microscope until cell layer is dispersed (usually within 5 to 15 minutes). Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal. Add 6.0 to 8.0 mL of complete growth medium and aspirate cells by gently pipetting. Add appropriate aliquots of the cell suspension to new culture vessels. Incubate cultures at 37°C. Population doubling time of approximately 17 hours. NOTE: For more information on enzymatic dissociation and subculturing of cell lines consult Chapter 12 in Culture of Animal Cells, a manual of Basic Technique by R. Ian Freshney, 6th edition, published by Alan R. Liss, N.Y., 2010.

Subculturing Medium Renewal:

Every 2 to 3 days

Culture Conditions:

Atmosphere: air, 95%; carbon dioxide (CO₂), 5% Temperature: 37°C

Cryopreservation:

95% FBS + 5% DMSO (Dimethyl sulfoxide)

Thawing Frozen Cells:

SAFETY PRECAUTION: It is highly recommended that protective gloves and clothing always be used and a full face mask always be worn when handling frozen vials. It is important to note that some vials leak when submerged in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vessel exploding or blowing off its cap with dangerous force creating flying debris. 1. Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 minutes). 2. Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions. 3. For cells that are sensitive to DMSO it is recommended that the cryoprotective agent be removed immediately. Transfer the vial contents to a centrifuge tube containing 9.0 mL complete culture medium and spin at approximately 125 x g for 5 to 7 minutes. 4. Discard the supernatant and Resuspend cell pellet with the recommended complete medium (see the specific batch information for the culture recommended dilution ratio). 5. Incubate the culture in an appropriate atmosphere and temperature (see "Culture Conditions" for this cell line). **NOTE:** It is important to avoid excessive alkalinity of the medium during recovery of the cells. It is suggested that, prior to the addition of the vial contents, the culture vessel containing the growth medium be placed into the incubator for at least 15 minutes to allow the medium to reach its normal pH (7.0 to 7.6).

References:

Huynh HT1, Robitaille G, Turner JD. Exp Cell Res. 1991 Dec;197(2):191-9. Rejman JJ1, Oliver SP, Muenchen RA, Turner JD. Cell Biol Int Rep. 1992 Oct;16(10):993-1001.

Depositors:

MÔNICA MARIA OLIVEIRA PINHO CERQUEIRA - Universidade Federal de Minas Gerais

ATCC:

CRL-10274