

Data Sheet

PAGE 1/4

| | |
|---------------------------|---|
| BCRJ Code: | 0164 |
| Cell Line: | MDA-MB-231 |
| Species: | Homo sapiens |
| Vulgar Name: | Human |
| Tissue: | Mammary Gland/Breast; Derived From Metastatic Site: Pleural Effusion |
| Cell Type: | Epithelial |
| Morphology: | Epithelial |
| Disease: | Adenocarcinoma |
| Growth Properties: | Adherent |
| Sex: | Female |
| Age/Ethnicity: | 51 Year / Caucasian |
| Applications: | These cells are a suitable transfection host |
| DNA Profile: | Amelogenin: X CSF1PO: 12,13 D13S317: 13 D16S539: 12 D5S818: 12 D7S820: 8,9 TH01: 7,9.3 TPOX: 8,9 vWA: 15,18 |
| Tumor Formation:: | Yes, in ALS treated BALB/c mice, forms poorly differentiated adenocarcinoma (grade III) Yes, in nude mice, forms poorly differentiated adenocarcinoma (grade III) |
| Biosafety: | 1 |
| Culture Medium: | Leibovitz's L-15 Medium contains 2 mM L-glutamine, NO sodium bicarbonate and fetal bovine serum to a final concentration of 10%. Note: The L-15 medium formulation was devised for use in a free gas exchange with atmospheric air. A CO ₂ and air mixture is detrimental to cells when using this medium for cultivation. |

Data Sheet

PAGE 2/4

Subculturing:

Volumes used in this protocol are for 75 cm² flasks; proportionally reduce or increase amount of dissociation medium for culture vessels of other sizes. Remove and discard culture medium. Briefly rinse the cell layer with PBS without calcium and magnesium to remove all traces of serum which contains trypsin inhibitor. Add 2.0 to 3.0 mL of Trypsin-EDTA solution to flask and observe cells under an inverted microscope until cell layer is dispersed (usually within 5 to 15 minutes). Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal. Add 6.0 to 8.0 mL of complete growth medium and aspirate cells by gently pipetting. Add appropriate aliquots of the cell suspension to new culture vessels. Incubate cultures at 37°C. NOTE: For more information on enzymatic dissociation and subculturing of cell lines consult Chapter 12 in Culture of Animal Cells, a manual of Basic Technique by R. Ian Freshney, 6th edition, published by Alan R. Liss, N.Y., 2010.

Subculturing Medium Renewal:

2 to 3 times per week

Subculturing Subcultivation Ratio:

1:2 to 1:4

Culture Conditions:

Atmosphere: air, 100% Temperature: 37°C

Cryopreservation:

95% FBS + 5% DMSO (Dimethyl sulfoxide)

Thawing Frozen Cells:

SAFETY PRECAUTION: It is highly recommended that protective gloves and clothing always be used and a full face mask always be worn when handling frozen vials. It is important to note that some vials leak when submerged in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vessel exploding or blowing off its cap with dangerous force creating flying debris.

1. Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the vial and cap out of the water. Thawing should be rapid (approximately 2 minutes).
2. Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
3. For cells that are sensitive to DMSO it is recommended that the cryoprotective agent be removed immediately. Transfer the vial contents to a centrifuge tube containing 9.0 mL complete culture medium and spin at approximately 125 x g for 5 to 7 minutes.
4. Discard the supernatant and Resuspend cell pellet with the recommended complete medium (see the specific batch information for the culture recommended dilution ratio).
5. Incubate the culture in a appropriate atmosphere and temperature (see "Culture Conditions" for this cell line).

NOTE: It is important to avoid excessive alkalinity of the medium during recovery of the cells. It is suggested that, prior to the addition of the vial contents, the culture vessel containing the growth medium be placed into the incubator for at least 15 minutes to allow the medium to reach its normal pH (7.0 to 7.6).

Data Sheet

PAGE 4/4

References:

Brinkley BR, et al. Variations in cell form and cytoskeleton in human breast carcinoma cells in vitro. *Cancer Res.* 40: 3118-3129, 1980. PubMed: 7000337

Cruciger Q, et al. Morphological, biochemical and chromosomal characterization of breast tumor lines from pleural effusions. *In Vitro* 12: 331, 1976. Siciliano MJ, et al. Mutually exclusive genetic signatures of human breast tumor cell lines with a common chromosomal marker. *Cancer Res.* 39: 919-922, 1979. PubMed: 427779

Cailleau R, et al. Breast tumor cell lines from pleural effusions. *J. Natl. Cancer Inst.* 53: 661-674, 1974. PubMed: 4412247

Cailleau R, et al. Long-term human breast carcinoma cell lines of metastatic origin: preliminary characterization. *In Vitro* 14: 911-915, 1978. PubMed: 730202

Bates SE, et al. Expression of the transforming growth factor-alpha/epidermal growth factor receptor pathway in normal human breast epithelial cells. *Endocrinology* 126: 596-607, 1990. PubMed: 2294006

Dickstein B, et al. Increased epidermal growth factor receptor in an estrogen-responsive, adriamycin-resistant MCF-7 cell line. *J. Cell. Physiol.* 157: 110-118, 1993. PubMed: 8408230

Huguet EL, et al. Differential expression of human Wnt genes 2, 3, 4, and 7B in human breast cell lines and normal and disease states of human breast tissue. *Cancer Res.* 54: 2615-2621, 1994. PubMed: 8168088

Satya-Prakash KL, et al. Cytogenetic analysis on eight human breast tumor cell lines: high frequencies of 1q, 11q and HeLa-like marker chromosomes. *Cancer Genet. Cytogenet.* 3: 61-73, 1981. PubMed: 7272986

Katayose Y, et al. Promoting apoptosis: a novel activity associated with the Cyclin-dependent kinase inhibitor p27. *Cancer Res.* 57: 5441-5445, 1997. PubMed: 9407946

Littlewood-Evans AJ, et al. The osteoclast-associated protease cathepsin K is expressed in human breast carcinoma. *Cancer Res.* 57: 5386-5390, 1997. PubMed: 9393764

Sheng S, et al. Maspin acts at the cell membrane to inhibit invasion and motility of mammary and prostatic cancer cells. *Proc. Natl. Acad. Sci. USA* 93: 11669-11674, 1996. PubMed: 8876194

De Vincenzo R, et al. Antiproliferative activity of colchicine analogues on MDR-positive and MDR-negative human cancer cell lines. *Anticancer Drug Des.* 13: 19-33, 1998. PubMed: 9474240

Soker S, et al. Characterization of novel vascular endothelial growth factor (VEGF) receptors on tumor cells that bind VEGF165 via its exon 7-encoded domain. *J. Biol. Chem.* 271: 5761-5767, 1996. PubMed: 8621443

Huguet EL, et al. Differential expression of human Wnt genes 2, 3, 4, and 7B in human breast cell lines and normal and disease states of human breast tissue. *Cancer Res.* 54: 2615-2621, 1994. PubMed: 8168088

Depositors:

RUY GASTALDONI JAEGER; Universidade de São Paulo.

ATCC:

HTB-26