

Data Sheet

PAGE 1/4

BCRJ Code:	0180
Cell Line:	MRC-5
Species:	Homo sapiens
Vulgar Name:	Human
Tissue:	Lung
Cell Type:	Fibroblast
Morphology:	Fibroblast
Disease:	Normal
Growth Properties:	Adherent
Sex:	Male
Age/Ethnicity:	14 (gestation) Week / Caucasian
Derivation:	The MRC-5 cell line was derived from normal lung tissue of a 14 week-old male fetus.
Applications:	efficacy testing transfection host viruscide testing
DNA Profile:	Amelogenin: X,Y CSF1PO: 11,12 D13S317: 11,14 D16S539: 9,11 D5S818: 11,12 D7S820: 10,11 THO1: 8 TPOX: 8 vWA: 15
Virus Susceptibility::	Human poliovirus 1 Herpes simplex virus Vesicular stomatitis, Glasgow (Indiana) Vesicular stomatitis, Orsay (Indiana)
Biosafety:	1
Additional Info:	The cells are capable of attaining 42-46 populatio doubling before senescence.



Data Sheet

PAGE 2/4

Culture Medium:

Dulbecco's Modified Eagle's Medium (DMEM) with 1% non-essential amino acids, 2 mM L-glutamine, 1 mM sodium pyruvate, 1.0 g/L glucose and 10% of fetal bovine serum.

Subculturing:

Volumes used in this protocol are for 75 cm² flask; proportionally reduce or increase amount of dissociation medium for culture vessels of other sizes. Remove and discard culture medium. Briefly rinse the cell layer with PBS without calcium and magnesium to remove all traces of serum which contains trypsin inhibitor. Add 2.0 to 3.0 mL of Trypsin-EDTA solution to flask and observe cells under an inverted microscope until cell layer is dispersed (usually within 5 to 15 minutes). Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal. Add 6.0 to 8.0 mL of complete growth medium and aspirate cells by gently pipetting. Add appropriate aliquots of the cell suspension to new culture vessels. Incubate cultures at 37°C. NOTE: For more information on enzymatic dissociation and subculturing of cell lines consult Chapter 12 in Culture of Animal Cells, a manual of Basic Technique by R. Ian Freshney, 6th edition, published by Alan R. Liss, N.Y., 2010.

Subculturing Medium Renewal:

1 to 2 times per week

Subculturing Subcultivation Ratio:

1:2 to 1:5

Culture Conditions:

Atmosphere: air, 95%; carbon dioxide (CO₂), 5% Temperature: 37°C

Cryopreservation:

95% FBS + 5% DMSO (Dimethyl sulfoxide)

SAFETY PRECAUTION: It is strongly recommended to always wear protective gloves, clothing, and a full-face mask when handling frozen vials. Some vials may leak when submerged in liquid nitrogen, allowing nitrogen to slowly enter the vial. Upon thawing, the conversion of liquid nitrogen back to its gas phase may cause the vial to explode or eject its cap with significant force, creating flying debris.

1. Thaw the vial by gently agitating it in a 37°C water bath. To minimize contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 minutes).

2. Remove the vial from the water bath as soon as its contents are thawed and decontaminate it by dipping in or spraying with 70% ethanol. From this point, all operations must be performed under strict aseptic conditions.

3. For cells sensitive to DMSO, it is recommended to remove the cryoprotective agent immediately. Transfer the vial contents to a centrifuge tube containing 9.0 mL of complete culture medium and centrifuge at approximately 125 × g for 5 to 7 minutes.

4. Discard the supernatant and resuspend the cell pellet in the recommended complete medium (see specific batch information for the appropriate dilution ratio).

5. Incubate the culture under appropriate atmospheric and temperature conditions (see "Culture Conditions" for this cell line).

NOTE: It is important to avoid excessive alkalinity of the medium during cell recovery. To minimize this risk, it is recommended to place the culture vessel containing the growth medium in the incubator for at least 15 minutes before adding the vial contents. This allows the medium to stabilize at its normal pH (7.0 to 7.6).

Jacobs JP, et al. Characteristics of a human diploid cell designated MRC-5. *Nature* 227: 168-170, 1970. PubMed: 4316953 . Proc. Symp. Human Diploid Cells. Zagreb, Yugoslavia: Yugoslav Acad. Sci. Arts; 1970. Holliday R, Tarrant GM. Altered enzymes in ageing human fibroblasts. *Nature* 238: 26-30, 1972. PubMed: 12635262 Lewis CM, Tarrant GM. Error theory and ageing in human diploid fibroblasts. *Nature* 239: 316-318, 1972. PubMed: 12635218 Landers JE, et al. Translational enhancement of mdm2 oncogene expression in human tumor cells containing a stabilized wild-type p53 protein. *Cancer Res.* 57: 3562-3568, 1997. PubMed: 9270029 Kaur A, et al. Cytotoxic T-lymphocyte responses to cytomegalovirus in normal and simian immunodeficiency virus-infected Rhesus macaques. *J. Virol.* 70: 7725-7733, 1996. PubMed: 8892893

Thawing Frozen Cells:

References:

Depositors:

José Mauro Granjeiro - INMETRO

Cellosaurus:

[CVCL_0440](#)



