

## Data Sheet

<b>BCRJ Code:</b>	0352
<b>Cell Line:</b>	NCI-H1975 [H-1975, H1975]
<b>Species:</b>	Homo sapiens
<b>Vulgar Name:</b>	Human
<b>Tissue:</b>	Lung
<b>Morphology:</b>	Epithelial
<b>Disease:</b>	Adenocarcinoma, Non-Small Cell Lung Cancer
<b>Growth Properties:</b>	Adherent
<b>Sex:</b>	Female
<b>Derivation:</b>	The line was established in July 1988. The tissue donor was a non-smoker.
<b>Biosafety:</b>	1
<b>Culture Medium:</b>	RPMI-1640 medium modified to contain 2 mM L-glutamine, 4500 mg/L glucose with fetal bovine serum to a final concentration of 10%.
<b>Subculturing:</b>	Remove medium, and rinse the monolayer with PBS without calcium and magnesium. Remove the trypsin, add fresh trypsin and let the culture sit at room temperature (or at 37°C) until the cells detach (about 10 minutes). Add fresh medium, aspirate and dispense into new flasks. T-75 flasks are recommended for subculturing this product. NOTE: For more information on enzymatic dissociation and subculturing of cell lines consult Chapter 12 in Culture of Animal Cells, a manual of Basic Technique by R. Ian Freshney, 6th edition, published by Alan R. Liss, N.Y., 2010.
<b>Subculturing Medium Renewal:</b>	Two to three times weekly

**Data Sheet**

PAGE 2/3

**Subculturing  
Subcultivation Ratio:**

1:3 to 1:6

**Culture Conditions:**Atmosphere: air, 95%; carbon dioxide (CO<sub>2</sub>), 5% Temperature: 37°C**Cryopreservation:**

95% FBS + 5% DMSO (Dimethyl sulfoxide)

**Thawing Frozen Cells:**

**SAFETY PRECAUTION:** It is highly recommended that protective gloves and clothing always be used and a full face mask always be worn when handling frozen vials. It is important to note that some vials leak when submerged in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vessel exploding or blowing off its cap with dangerous force creating flying debris. 1. Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 minutes). 2. Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions. 3. For cells that are sensitive to DMSO it is recommended that the cryoprotective agent be removed immediately. Transfer the vial contents to a centrifuge tube containing 9.0 mL complete culture medium and spin at approximately 125 x g for 5 to 7 minutes. 4. Discard the supernatant and Resuspend cell pellet with the recommended complete medium (see the specific batch information for the culture recommended dilution ratio). 5. Incubate the culture in an appropriate atmosphere and temperature (see "Culture Conditions" for this cell line). **NOTE:** It is important to avoid excessive alkalinity of the medium during recovery of the cells. It is suggested that, prior to the addition of the vial contents, the culture vessel containing the growth medium be placed into the incubator for at least 15 minutes to allow the medium to reach its normal pH (7.0 to 7.6).

**References:**

NCI-Navy Medical Oncology Branch Cell Line Supplement. J. Cell. Biochem. suppl. 24: 1996. Sordella R, et al. Gefitinib-sensitizing EGFR mutations in lung cancer activate anti-apoptotic pathways. Science 305: 1163-1167, 2004. PubMed: 15284455

**Depositors:**

Lidia Moreira Lima - UNIVERSIDADE FEDERAL DO RIO DE JANEIRO Rui Manoel Reis - Fundação Pio XII

**ATCC:**

CRL-5908

**Cellosaurus:**[CVCL 1511](#)

