

Data Sheet

BCRJ Code:	0189
Cell Line:	Neuro-2a
Species:	Mus musculus
Vulgar Name:	Mouse; A Albino
Tissue:	Brain
Cell Type:	Neuroblast
Morphology:	Neuronal And Amoeboid Stem Cells
Disease:	Neuroblastoma
Growth Properties:	Adherent
Derivation:	Derived from a spontaneous tumour in an albino strain A mouse. Cells produce microtubular protein which is believed to play a role in the contractile system giving axoplasmic flow in nerve cells.
Applications:	This cell line is a suitable transfection host. The cell line has been used for studies on the mechanism of vinblastine precipitation of microtubular protein, the kinetics of GTP binding to isolated protein, the turnover of microtubules in vivo, and the synthesis and assembly of microtubular protein . The World Organization for Animal Health (OIE) uses the cells for routine diagnosis of rabies.
Virus Susceptibility::	Herpes simplex virus Vesicular stomatitis virus Human poliovirus 1
Products:	acetylcholinesterase; tubulin
Biosafety:	1
Culture Medium:	Dulbecco's Modified Eagle's Medium (DMEM) with 1% non-essential amino acids, 2 mM L-glutamine, 4.5 g/L glucose and 10% of fetal bovine serum, 10%.

Subculturing:

Volumes used in this protocol are for 75 cm² flask; proportionally reduce or increase amount of dissociation medium for culture vessels of other sizes. Remove and discard culture medium. Briefly rinse the cell layer with PBS without calcium and magnesium to remove all traces of serum which contains trypsin inhibitor. Add 2.0 to 3.0 mL of Trypsin-EDTA solution to flask and observe cells under an inverted microscope until cell layer is dispersed (usually within 5 to 15 minutes). Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal. Add 6.0 to 8.0 mL of complete growth medium and aspirate cells by gently pipetting. Add appropriate aliquots of the cell suspension to new culture vessels. Incubate cultures at 37°C. NOTE: For more information on enzymatic dissociation and subculturing of cell lines consult Chapter 12 in Culture of Animal Cells, a manual of Basic Technique by R. Ian Freshney, 6th edition, published by Alan R. Liss, N.Y., 2010.

Subculturing Medium Renewal:

1 to 2 times per week

Subculturing Subcultivation Ratio:

1:3 to 1:6

Culture Conditions:

Atmosphere: air, 95%; carbon dioxide (CO₂), 5% Temperature: 37°C

Cryopreservation:

95% FBS + 5% DMSO (Dimethyl sulfoxide)

Thawing Frozen Cells:

SAFETY PRECAUTION: It is highly recommended that protective gloves and clothing always be used and a full face mask always be worn when handling frozen vials. It is important to note that some vials leak when submerged in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vessel exploding or blowing off its cap with dangerous force creating flying debris. 1. Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the Oring and cap out of the water. Thawing should be rapid (approximately 2 minutes). 2. Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions. 3. For cells that are sensitive to DMSO it is recommended that the cryoprotective agent be removed immediately. Transfer the vial contents to a centrifuge tube containing 9.0 mL complete culture medium and spin at approximately 125 x g for 5 to 7 minutes. 4. Discard the supernatant and Resuspend cell pellet with the recommended complete medium (see the specific batch information for the culture recommended dilution ratio). 5. Incubate the culture in an appropriate atmosphere and temperature (see "Culture Conditions" for this cell line). **NOTE:** It is important to avoid excessive alkalinity of the medium during recovery of the cells. It is suggested that, prior to the addition of the vial contents, the culture vessel containing the growth medium be placed into the incubator for at least 15 minutes to allow the medium to reach its normal pH (7.0 to 7.6).

References:

Olmsted JB, et al. Isolation of microtubule protein from cultured mouse neuroblastoma cells. Proc. Natl. Acad. Sci. USA 65: 129-136, 1970. PubMed: 5263744 Klebe RJ, Ruddle FH. Neuroblastoma: Cell culture analysis of a differentiating stem cell system. J. Cell Biol. 43: 69A, 1969. Naslavsky N, et al. Characterization of detergent-insoluble complexes containing the cellular prion protein and its scrapie isoform. J. Biol. Chem. 272: 6324-6331, 1997. PubMed: 9045652 Kaneko K, et al. Evidence for protein X binding to a discontinuous epitope on the cellular prion protein during scrapie prion propagation. Proc. Natl. Acad. Sci. USA 94: 10069-10074, 1997. PubMed: 9294164 Maestrini E, et al. A family of transmembrane proteins with homology to the MET-hepatocyte growth factor receptor. Proc. Natl. Acad. Sci. USA 93: 674-678, 1996. PubMed: 8570614

Depositors:

Elizabeth Obino Lima, Fundação Instituto Oswaldo Cruz.

ATCC:

CCL-131

Cellosaurus:

[CVCL_0470](#)

