

Data Sheet

PAGE 1/3

BCRJ Code:	0294
Cell Line:	ONS-76
Species:	Homo sapiens
Vulgar Name:	Human
Tissue:	Neural/Brain
Cell Type:	Epithelial
Morphology:	Epithelial-Like
Disease:	Medulloblastoma
Growth Properties:	Adherent
Sex:	Female
Age/Ethnicity:	2 Year /
DNA Profile:	AMELOGENINA: X D5S818: 9,10 D13S317: 8,13 D7S820: 11,12 D16S539: 9,10 vWA: 14,16 TH01: 9 TPOX: 8 CSF1PO:10,12
Biosafety:	1
Additional Info:	POSITIVE FOR 145 Kda AND 200kDa NEUROFILAMENT AND NSE. Negative for GFAP AND S-100 PROTEIN
Culture Medium:	RPMI-1640 medium modified to contain 2 mM L-glutamine, 4500 mg/L glucose and 10% of fetal bovine serum.

Subculturing:

Remove and discard culture medium. Briefly rinse the cell layer with PBS without calcium and magnesium to remove all traces of serum which contains trypsin inhibitor. Add 2.0 to 3.0 mL of Trypsin-EDTA solution to flask and observe cells under an inverted microscope until cell layer is dispersed (usually within 5 to 15 minutes). Add 6.0 to 8.0 mL of complete growth medium and aspirate cells by gently pipetting. Add appropriate aliquots of the cell suspension into new culture vessels. Incubate cultures at 37°C. NOTE: For more information on enzymatic dissociation and subculturing of cell lines consult Chapter 12 in Culture of Animal Cells, a manual of Basic Technique by R. Ian Freshney, 6th edition, published by Alan R. Liss, N.Y., 2010.

Subculturing Medium Renewal:

2 to 3 times per week

Culture Conditions:

Atmosphere: air, 95%; carbon dioxide (CO₂), 5% Temperature: 37°C

Cryopreservation:

95% FBS + 5% DMSO (Dimethyl sulfoxide)

Thawing Frozen Cells:

SAFETY PRECAUTION: It is highly recommended that protective gloves and clothing always be used and a full face mask always be worn when handling frozen vials. It is important to note that some vials leak when submerged in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vessel exploding or blowing off its cap with dangerous force creating flying debris. 1. Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 minutes). 2. Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions. 3. For cells that are sensitive to DMSO is recommended that the cryoprotective agent be removed immediately. Transfer the vial contents to a centrifuge tube containing 9.0 mL complete culture medium and spin at approximately 125 x g for 5 to 7 minutes. 4. Discard the supernatant and Resuspend cell pellet with the recommended complete medium (see the specific batch information for the culture recommended dilution ratio). 5. Incubate the culture in an appropriate atmosphere and temperature (see "Culture Conditions" for this cell line). NOTE: It is important to avoid excessive alkalinity of the medium during recovery of the cells. It is suggested that, prior to the addition of the vial contents, the culture vessel containing the growth medium be placed into the incubator for at least 15 minutes to allow the medium to reach its normal pH (7.0 to 7.6).

References:

.Shimizu,K., Yamada,M., Moriuchi,S., Tamura,K., Okamoto,Y., Park,K.-C., Mabuchi,E., Miyao,Y., and Hayakawa,T. Human medulloblastoma cell lines: ONS-76 and ONS-81, Hum. Cell, 3: 265-266, 1990. Tamura,K., Shimizu,K., Yamada,M., Okamoto,Y., Matsui,Y., Park,K.-C., Mabuchi,E., Moriuchi,S., and Mogami,H. Expression of Major histocompatibility complex on human medulloblastoma cells with neuronal differentiation, Cancer Res., 49: 5380-5384, 1989.

Depositors:

Luíz Gonzaga Tone; Universidade de São Paulo

Cellosaurus:

[CVCL_1624](#)