

Data Sheet

BCRJ Code:	0200
Cell Line:	P815
Species:	Mus musculus
Vulgar Name:	Mouse; Dbal/2
Tissue:	Mast Cell
Disease:	Mastocytoma
Growth Properties:	Suspension (Some Adherent Cells)
Derivation:	established from the mastocytoma tumor of a DBA/2 mouse treated with methylcolanthrene; used as target cells for cytotoxic T cell assays; as reported cells exhibit no effector activity in an antibody-dependent cell mediated cytotoxic system
Applications:	This cell line is a suitable transfection host.
Products:	Lysozyme
Biosafety:	1
Additional Info:	P815 cells phagocytose latex beads but not zymosan or BCG. They do not function in antibody dependent cell mediated cytotoxicity. Growth of the cells is not inhibited by dextran sulfate, LPS or PPD.
Culture Medium:	Dulbecco's Modified Eagle's Medium (DMEM) modified to contain 2 mM L-glutamine, 4500 mg/L glucose and 10% of fetal bovine serum.
Subculturing:	Cultures can be maintained by addition or replacement of fresh medium. Start cultures at 2 X 10 ⁵ cells/mL and maintain between 1 X 10 ⁵ and 1 X 10 ⁶ cells/mL. Adherent cells can be recovered by scraping. NOTE: Maximum cell density at > 1 x 10 ⁶ cells/ml Population Doubling Time: 18-22 hrs

**Subculturing Medium
Renewal:**

Every 2 to 3 days

Culture Conditions:

Atmosphere: air, 95%; carbon dioxide (CO₂), 5% Temperature: 37°C

Cryopreservation:

95% FBS + 5% DMSO (Dimethyl sulfoxide)

Thawing Frozen Cells:

SAFETY PRECAUTION: It is highly recommended that protective gloves and clothing always be used and a full face mask always be worn when handling frozen vials. It is important to note that some vials leak when submerged in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vessel exploding or blowing off its cap with dangerous force creating flying debris. 1. Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 minutes). 2. Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions. 3. For cells that are sensitive to DMSO it is recommended that the cryoprotective agent be removed immediately. Transfer the vial contents to a centrifuge tube containing 9.0 mL complete culture medium and spin at approximately 125 x g for 5 to 7 minutes. 4. Discard the supernatant and Resuspend cell pellet with the recommended complete medium (see the specific batch information for the culture recommended dilution ratio). 5. Incubate the culture in an appropriate atmosphere and temperature (see "Culture Conditions" for this cell line). **NOTE:** It is important to avoid excessive alkalinity of the medium during recovery of the cells. It is suggested that, prior to the addition of the vial contents, the culture vessel containing the growth medium be placed into the incubator for at least 15 minutes to allow the medium to reach its normal pH (7.0 to 7.6).

References:

Ralph P, et al. Lysozyme synthesis by established human and murine histiocytic lymphoma cell lines. *J. Exp. Med.* 143: 1528-1533, 1976. PubMed: 1083890 Ralph P, Nakoinz I. Antibody-dependent killing of erythrocyte and tumor targets by macrophage-related cell lines: enhancement by PPD and LPS. *J. Immunol.* 119: 950-954, 1977. PubMed: 894031 Ralph P, Nakoinz I. Direct toxic effects of immunopotentiators on monocytic myelomonocytic, and histiocytic or macrophage tumor cells in culture. *Cancer Res.* 37: 546-550, 1977. PubMed: 318922 Ralph P, Nakoinz I. Lipopolysaccharides inhibit lymphosarcoma cells of bone marrow origin. *Nature* 249: 49-51, 1974. PubMed: 4208429 Ralph P, et al. Lymphosarcoma cell growth is selectively inhibited by B lymphocyte mitogens: LPS, dextran sulfate and PPD. *Biochem. Biophys. Res. Commun.* 61: 1268-1275, 1974. PubMed: 4616699 Lundak RL, Raidt DJ. Cellular immune response against tumor cells. I. In vitro immunization of allogeneic and syngeneic mouse spleen cell suspensions against DBA mastocytoma cells. *Cell. Immunol.* 9: 60-66, 1973. PubMed: 4270287 Plaut M, et al. Studies on the mechanism of lymphocyte-mediated cytotoxicity. IV. Specificity of the histamine receptor on effector T cells. *J. Immunol.* 111: 389-394, 1973. PubMed: 4123976 Schmidt W, et al. Cell-free tumor antigen peptide-based cancer vaccines. *Proc. Natl. Acad. Sci. USA* 94: 3262-3267, 1997. PubMed: 9096381 Gonzalez Armas JC, et al. DNA immunization confers protection against murine cytomegalovirus infection. *J. Virol.* 70: 7921-7928, 1996. PubMed: 8892915

Depositors:

Dr. T. kipnis, Universidade de São Paulo.

ATCC:

TIB-64