

Data Sheet

BCRJ Code:	0269
Cell Line:	PC-3
Species:	Homo sapiens
Vulgar Name:	Human
Tissue:	Prostate; Derived From Metastatic Site: Bone
Morphology:	Epithelial
Disease:	Adenocarcinoma (Grade Iv)
Growth Properties:	Adherent (The Cells Form Clusters In Soft Agar And Can Be Adapted To Suspension Growth)
Sex:	Male
Age/Ethnicity:	62 Year / Caucasian
Derivation:	The PC-3 was initiated from a bone metastasis of a grade IV prostatic adenocarcinoma from a 62-year-old male Caucasian.
Applications:	This cell line is a suitable transfection host.
DNA Profile:	Amelogenin: X CSF1PO: 11 D13S317: 11 D16S539: 11 D5S818: 13 D7S820: 8,11 THO1: 6,7 TPOX: 8,9 vWA: 17
Tumor Formation::	Yes, in semi-solid medium Yes, tumors developed within 21 days at 100% frequency (5/5) in nude mice inoculated subcutaneously with 10(7) cells
Biosafety:	1
Additional Info:	The cells exhibit low acid phosphatase and testosterone-5-alpha reductase activities.

Culture Medium:

F-12K Medium (Kaighn's Modification of Ham's F-12 Medium) contains 2 mM L-glutamine and 10% of fetal bovine serum.

Subculturing:

Volumes used in this protocol are for 75 cm² flask; proportionally reduce or increase amount of dissociation medium for culture vessels of other sizes. Remove and discard culture medium. Briefly rinse the cell layer with PBS without calcium and magnesium to remove all traces of serum that contains trypsin inhibitor. Add 2.0 to 3.0 mL of Trypsin-EDTA solution to flask and observe cells under an inverted microscope until cell layer is dispersed (usually within 5 to 15 minutes). Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal. Add 6.0 to 8.0 mL of complete growth medium and aspirate cells by gently pipetting. Add appropriate aliquots of the cell suspension to new culture vessels. Incubate cultures at 37°C. NOTE: For more information on enzymatic dissociation and subculturing of cell lines consult Chapter 12 in Culture of Animal Cells, a manual of Basic Technique by R. Ian Freshney, 6th edition, published by Alan R. Liss, N.Y., 2010.

Subculturing Medium Renewal:

2 to 3 times a week

Subculturing Subcultivation Ratio:

1:3 to 1:6 is recommended

Culture Conditions:

Atmosphere: air, 95%; carbon dioxide (CO₂), 5% Temperature: 37°C

Cryopreservation:

95% FBS + 5% DMSO (Dimethyl sulfoxide)

Thawing Frozen Cells:

SAFETY PRECAUTION: It is highly recommended that protective gloves and clothing always be used and a full face mask always be worn when handling frozen vials. It is important to note that some vials leak when submerged in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vessel exploding or blowing off its cap with dangerous force creating flying debris.

1. Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 minutes).
2. Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
3. For cells that are sensitive to DMSO it is recommended that the cryoprotective agent be removed immediately. Transfer the vial contents to a centrifuge tube containing 9.0 mL complete culture medium and spin at approximately 125 x g for 5 to 7 minutes.
4. Discard the supernatant and Resuspend cell pellet with the recommended complete medium (see the specific batch information for the culture recommended dilution ratio).
5. Incubate the culture in an appropriate atmosphere and temperature (see "Culture Conditions" for this cell line).

NOTE: It is important to avoid excessive alkalinity of the medium during recovery of the cells. It is suggested that, prior to the addition of the vial contents, the culture vessel containing the growth medium be placed into the incubator for at least 15 minutes to allow the medium to reach its normal pH (7.0 to 7.6).

References:

22363: Kaighn ME, et al. Establishment and characterization of a human prostatic carcinoma cell line (PC-3). Invest. Urol. 17: 16-23, 1979. PubMed: 447482 22470: Chen TR. Chromosome identity of human prostate cancer cell lines, PC-3 and PPC-1. Cytogenet. Cell Genet. 62: 183-184, 1993. PubMed: 8428522 26302: Ohnuki Y, et al. Chromosomal analysis of human prostatic adenocarcinoma cell lines. Cancer Res. 40: 524-534, 1980. PubMed: 7471073 32341: Sheng S, et al. Maspin acts at the cell membrane to inhibit invasion and motility of mammary and prostatic cancer cells. Proc. Natl. Acad. Sci. USA 93: 11669-11674, 1996. PubMed: 8876194 32344: Umekita Y, et al. Human prostate tumor growth in athymic mice: inhibition by androgens and stimulation by finasteride. Proc. Natl. Acad. Sci. USA 93: 11802-11807, 1996. PubMed: 8876218 32460: Carter RE, et al. Prostate-specific membrane antigen is a hydrolase with substrate and pharmacologic characteristics of a neuropeptidase. Proc. Natl. Acad. Sci. USA 93: 749-753, 1996. PubMed: 8570628 32486: Nupponen NN, et al. Genetic alterations in prostate cancer cell lines detected by comparative genomic hybridization. Cancer Genet. Cytogenet. 101: 53-57, 1998. PubMed: 9460501 32488: Geiger T, et al. Antitumor activity of a PKC-alpha antisense oligonucleotide in combination with standard chemotherapeutic agents against various human tumors transplanted into nude mice. Anticancer Drug Des. 13: 35-45, 1998. PubMed: 9474241 32916: Su ZZ, et al. Surface-epitope masking and expression cloning identifies the human prostate carcinoma tumor antigen gene PCTA-1 a member of the galectin gene family. Proc. Natl. Acad. Sci. USA 93: 7252-7257, 1996. PubMed: 8692978

Depositors:

Hernandes Faustino De Carvalho; Universidade Estadual De Campinas

ATCC:

CRL-1435

Cellosaurus:

[CVCL_0035](https://www.ebi.ac.uk/bioproject/100000/CVCL_0035)