

Data Sheet

PAGE 1/3

BCRJ Code:	0209
Cell Line:	R4-6A2
Species:	Rattus norvegicus (B cell); Mus musculus (myeloma), rat (B cell); mouse (myeloma)
Vulgar Name:	Rat; Da Atrud Institute
Cell Type:	Hybridoma: B Lymphocyte
Morphology:	Lymphoblast
Growth Properties:	Suspension
Derivation:	Animals were immunized with partially purified murine gamma interferon. Spleen cells were fused with P3U-1 myeloma cells.
Products:	Immunoglobulin; monoclonal antibody; gamma interferon: ab against; IgG1
Biosafety:	1
Additional Info:	The antibody neutralizes the ability of lymphokine preparations to induce tumoricidal activity in murine macrophages. The antibody neutralizes the antiviral effect of gamma interferon but does not affect the activities of murine alpha and beta interferon or rat or human gamma interferon. 30 ng of R4-6A2 antibody neutralizes the antiviral activity of 10 units of murine gamma interferon.
Culture Medium:	Dulbecco's Modified Eagle's Medium (DMEM) modified to contain 2 mM L-glutamine, 4500 mg/L glucose and 10% of fetal bovine serum.
Subculturing:	Cultures can be maintained by addition of fresh medium. Alternatively, cultures can be established by centrifugation with subsequent resuspension at 5×10^4 viable cells/mL. Maintain cultures at a cell concentration between 6×10^4 and 6×10^5 cells/mL. NOTE: Do not allow the cell concentration to reach 1×10^6 cells/mL.

Data Sheet

PAGE 2/3

Subculturing Medium Renewal:

Every 2 to 3 days

Culture Conditions:

Atmosphere: air, 95%; carbon dioxide (CO₂), 5% Temperature: 37°C

Cryopreservation:

95% FBS + 5% DMSO (Dimethyl sulfoxide)

SAFETY PRECAUTION: It is strongly recommended to always wear protective gloves, clothing, and a full-face mask when handling frozen vials. Some vials may leak when submerged in liquid nitrogen, allowing nitrogen to slowly enter the vial. Upon thawing, the conversion of liquid nitrogen back to its gas phase may cause the vial to explode or eject its cap with significant force, creating flying debris.

1. Thaw the vial by gently agitating it in a 37°C water bath. To minimize contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 minutes).
2. Remove the vial from the water bath as soon as its contents are thawed and decontaminate it by dipping in or spraying with 70% ethanol. From this point, all operations must be performed under strict aseptic conditions.
3. For cells sensitive to DMSO, it is recommended to remove the cryoprotective agent immediately. Transfer the vial contents to a centrifuge tube containing 9.0 mL of complete culture medium and centrifuge at approximately 125 × g for 5 to 7 minutes.
4. Discard the supernatant and resuspend the cell pellet in the recommended complete medium (see specific batch information for the appropriate dilution ratio).
5. Incubate the culture under appropriate atmospheric and temperature conditions (see "Culture Conditions" for this cell line).

Thawing Frozen Cells:

NOTE: It is important to avoid excessive alkalinity of the medium during cell recovery. To minimize this risk, it is recommended to place the culture vessel containing the growth medium in the incubator for at least 15 minutes before adding the vial contents. This allows the medium to stabilize at its normal pH (7.0 to 7.6).

References:

J. Exp. Med. 159: 1560-65, 1984.

Depositors:

Ises Abrahamsohn - Universidade de São Paulo

Cellosaurus:

[CVCL_9233](#)

