

Data Sheet

PAGE 1/3

| | |
|---------------------------|---|
| BCRJ Code: | 0267 |
| Cell Line: | RD-ES |
| Species: | Homo sapiens |
| Vulgar Name: | Human |
| Tissue: | Bone |
| Morphology: | Epithelial |
| Disease: | Ewing'S Sarcoma |
| Growth Properties: | Mixed: Adherent And Clusters In Suspension |
| Sex: | Male |
| Age/Ethnicity: | 19 Year / Caucasian |
| Derivation: | The cell line was initiated by G. Marshall and M. Kirchen from a primary osseous Ewings sarcoma of the humerus. |
| DNA Profile: | Amelogenin: X,Y CSF1PO:11 D13S317: 12,11 D16S539:11,9 D5S818:11 D7S820:10 THO1:7 TPOX:11,9 vWA:17 D2S1338:20,19 D19S433: 14,13 FGA:25,21 D3S1358:15 D18S51: 18,14 D8S1179: 13 D21S11: 28 |
| Products: | Antigen Expression: Blood type B; Rh+ |
| Biosafety: | 1 |
| Additional Info: | Ultrastructurally the cells exhibit primitive cell junctions, possess glycogen pools and are 20 to 25 microns in diameter. The cells grow as a loosely attached monolayer in small clusters of 5 to 10 cells. |
| Culture Medium: | RPMI-1640 medium modified to contain 2 mM L-glutamine, 4500 mg/L glucose and 10% of fetal bovine serum. |



Data Sheet

PAGE 2/3

Subculturing:

Shake the flask after removing most of the medium. Add fresh medium and transfer to new flasks.

Subculturing Medium Renewal:

2 to 3 times a week

Subculturing Subcultivation Ratio:

1:3 to 1:8 is recommended

Culture Conditions:

Atmosphere: air, 95%; carbon dioxide (CO₂), 5% Temperature: 37°C

Cryopreservation:

95% FBS + 5% DMSO (Dimethyl sulfoxide)

Thawing Frozen Cells:

SAFETY PRECAUTION: It is highly recommended that protective gloves and clothing always be used and a full face mask always be worn when handling frozen vials. It is important to note that some vials leak when submerged in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vessel exploding or blowing off its cap with dangerous force creating flying debris. 1. Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 minutes). 2. Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions. 3. For cells that are sensitive to DMSO it is recommended that the cryoprotective agent be removed immediately. Transfer the vial contents to a centrifuge tube containing 9.0 mL complete culture medium and spin at approximately 125 x g for 5 to 7 minutes. 4. Discard the supernatant and Resuspend cell pellet with the recommended complete medium (see the specific batch information for the culture recommended dilution ratio). 5. Incubate the culture in an appropriate atmosphere and temperature (see "Culture Conditions" for this cell line). NOTE: It is important to avoid excessive alkalinity of the medium during recovery of the cells. It is suggested that, prior to the addition of the vial contents, the culture vessel containing the growth medium be placed into the incubator for at least 15 minutes to allow the medium to reach its normal pH (7.0 to 7.6).

References:

Hay, R. J., Caputo, J.L., and Macy, M. L., Eds (1992), ATCC Quality Control Methods for Cell Lines. 2nd edition, Published by ATCC

Depositors:

Caroline Brunetto - Instituto do Câncer Infantil do RS



ATCC:

HTB-166