

## Data Sheet

PAGE 1/3

<b>BCRJ Code:</b>	0195
<b>Cell Line:</b>	SCC-4
<b>Species:</b>	Homo sapiens
<b>Vulgar Name:</b>	Human
<b>Tissue:</b>	Tongue
<b>Morphology:</b>	Epithelial-Like
<b>Disease:</b>	Squamous Cell Carcinoma
<b>Growth Properties:</b>	Adherent
<b>Sex:</b>	Male
<b>Age/Ethnicity:</b>	55 Year /
<b>DNA Profile:</b>	Amelogenin: X,Y CSF1PO: 11 D13S317: 11,13 D16S539: 12 D5S818: 13 D7S820: 9,11 TH01: 9.3 TPOX: 8 vWA: 15,17
<b>Tumor Formation::</b>	Yes, Tumors developed within 21 days at 100% frequency (5/5) in nude mice inoculated subcutaneously with 107 cells.
<b>Products:</b>	epidermal keratins; 40 kD keratin
<b>Biosafety:</b>	1
<b>Additional Info:</b>	SCC-4 forms colonies in semi-solid medium, and is not induced to differentiate by anchorage deprivation.
<b>Culture Medium:</b>	1:1 mixture of Dulbecco's modified Eagle's medium and Ham's F12 medium containing 1.2 g/L sodium bicarbonate, 2 mM L-glutamine, 400 ng/mL hydrocortisone and 10% of fetal bovine serum.

**Subculturing:**

Note: Subculture before 100% confluent. Volumes used in this protocol are for 75 cm<sup>2</sup> flasks; proportionally reduce or increase amount of dissociation medium for culture vessels of other sizes. Remove and discard culture medium. Briefly rinse the cell layer with 0.25% (w/v) Trypsin-0.53 mM EDTA solution to remove all traces of serum which contains trypsin inhibitor. Add 2.0 to 3.0 mL of Trypsin-EDTA solution to flask and observe cells under an inverted microscope until cell layer is dispersed (usually within 5 to 15 minutes). Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal. Add 6.0 to 8.0 mL of complete growth medium and aspirate cells by gently pipetting. Add appropriate aliquots of the cell suspension to new culture vessels. Inoculate new flasks with 3 x 10<sup>3</sup> cells/cm<sup>2</sup>. Incubate cultures at 37°C. Population Doubling Time: 60-70 hours NOTE: For more information on enzymatic dissociation and subculturing of cell lines consult Chapter 12 in Culture of Animal Cells, a manual of Basic Technique by R. Ian Freshney, 6th edition, published by Alan R. Liss, N.Y., 2010.

**Subculturing Medium Renewal:**

1 to 2 times per week

**Culture Conditions:**

Atmosphere: air, 95%; carbon dioxide (CO<sub>2</sub>), 5% Temperature: 37°C

**Cryopreservation:**

95% FBS + 5% DMSO (Dimethyl sulfoxide)

**Thawing Frozen Cells:**

**SAFETY PRECAUTION:** It is highly recommended that protective gloves and clothing always be used and a full face mask always be worn when handling frozen vials. It is important to note that some vials leak when submerged in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vessel exploding or blowing off its cap with dangerous force creating flying debris. 1. Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the Oring and cap out of the water. Thawing should be rapid (approximately 2 minutes). 2. Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions. 3. For cells that are sensitive to DMSO is recommended that the cryoprotective agent be removed immediately. Transfer the vial contents to a centrifuge tube containing 9.0 mL complete culture medium and spin at approximately 125 x g for 5 to 7 minutes. 4. Discard the supernatant and Resuspend cell pellet with the recommended complete medium (see the specific batch information for the culture recommended dilution ratio). 5. Incubate the culture in an appropriate atmosphere and temperature (see "Culture Conditions" for this cell line). **NOTE:** It is important to avoid excessive alkalinity of the medium during recovery of the cells. It is suggested that, prior to the addition of the vial contents, the culture vessel containing the growth medium be placed into the incubator for at least 15 minutes to allow the medium to reach its normal pH (7.0 to 7.6).

**References:**

Rheinwald JG, Beckett MA. Tumorigenic keratinocyte lines requiring anchorage and fibroblast support cultures from human squamous cell carcinomas. Cancer Res. 41: 1657-1663, 1981

**Depositors:**

ADELZON ASSIS DE PAULA; UFRJ

**ATCC:**

CRL-1624