

Banco de Células do Rio de Janeiro

Data Sheet

PAGE 1/3

BCRJ Code:	0344
Cell Line:	SJCRH30 [RC13, RMS 13, SJRH30]
Species:	Homo sapiens
Vulgar Name:	Human
Tissue:	Muscle, Derived From Metastatic Site: Bone Marrow
Cell Type:	Fibroblast
Morphology:	Fibroblast
Disease:	Rhabdomyosarcoma
Growth Properties:	Adherent
Sex:	Male
Age/Ethinicity:	17 Year / Caucasian
Derivation:	The line was established from cells from the bone marrow of a child with rhabdomyosarcoma. Derived from metastatic site, bone marrow
Products:	Genes Expressed: gli + (amplified 30 fold), N-myc + (not amplified)
Biosafety:	1
Addtional Info:	The cells show ultrastructural elements of primitive skeletal muscle differentiation.
Culture Medium:	RPMI-1640 medium modified to contain 2 mM L-glutamine, 4500 mg/L glucose and 10% of fetal bovine serum.

f

0



Banco de Células do Rio de Janeiro

Data Sheet

PAGE 2/3

Subculturing:	Remove medium, and rinse with PBS without calcium and magnesium . Remove the solution and add an additional 1 to 2 mL of trypsin-EDTA solution. Allow the flask to sit at room temperature (or at 37°C) until the cells detach. Add fresh culture medium, aspirate and dispense into new culture flasks. NOTE: For more information on enzymatic dissociation and subculturing of cell lines consult Chapter 12 in Culture of Animal Cells, a manual of Basic Technique by R. Ian Freshney, 6th edition, published by Alan R. Liss, N.Y., 2010.
Subculturing Medium Renewal:	Twice per week
Subculturing Subcultivation Ratio:	1:5 to 1:10 is recommended
Culture Conditions:	Atmosphere: air, 95%; carbon dioxide (CO2), 5% Temperature: 37°C
Cryopreservation:	95% FBS + 5% DMSO (Dimethyl sulfoxide)

f





Banco de Células do Rio de Janeiro

Data Sheet

PAGE 3/3

Thawing Frozen Cells:	SAFETY PRECAUTION: It is strongly recommended to always wear protective gloves, clothing, and a full-face mask when handling frozen vials. Some vials may leak when submerged in liquid nitrogen, allowing nitrogen to slowly enter the vial. Upon thawing, the conversion of liquid nitrogen back to its gas phase may cause the vial to explode or eject its cap with significant force, creating flying debris. 1. Thaw the vial by gently agitating it in a 37°C water bath. To minimize contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 minutes). 2. Remove the vial from the water bath as soon as its contents are thawed and decontaminate it by dipping in or spraying with 70% ethanol. From this point, all operations must be performed under strict aseptic conditions. 3. For cells sensitive to DMSO, it is recommended to remove the cryoprotective agent immediately. Transfer the vial contents to a centrifuge tube containing 9.0 mL of complete culture medium and centrifuge at approximately 125 × g for 5 to 7 minutes. 4. Discard the supernatant and resuspend the cell pellet in the recommended complete medium (see specific batch information for the appropriate dilution ratio). 5. Incubate the culture under appropriate atmospheric and temperature conditions (see "Culture Conditions" for this cell line). NOTE: It is important to avoid excessive alkalinity of the medium during cell recovery. To minimize this risk, it is recommended to place the culture vessel containing the growth medium in the incubator for at least 15 minutes before adding the vial contents. This allows the medium to
	stabilize at its normal pH (7.0 to 7.6).
References:	Oliner JD, et al. Amplification of a gene encoding a p53-associated protein in human sarcomas. Nature 358: 80-83, 1992. PubMed: 1614537 Roberts WM, et al. Amplification of the gli gene in childhood sarcomas. Cancer Res. 49: 5407-5413, 1989. PubMed: 2766305 Douglass EC, et al. A specific chromosomal abnormality in rhabdomyosarcoma [published erratum appears in Cytogenet Cell Genet 1988;47(4):following 232]. Cytogenet. Cell Genet. 45: 148-155, 1987. PubMed: 3691179
Depositors:	Magnus Regios Dias da Silva - UNIFESP

f

