

Data Sheet

PAGE 1/3

BCRJ Code:	0409
Cell Line:	SK-OV-3
Species:	Homo sapiens
Vulgar Name:	Human
Tissue:	Ovary
Cell Type:	Epithelial
Morphology:	Epithelial
Disease:	Adenocarcinoma
Growth Properties:	Adherent
Sex:	Female
Age/Ethnicity:	64 Year /
Derivation:	Derived from the ascitic fluid from a 64 year old caucasian female with an ovarian tumour. Forms moderately well-differentiated adenocarcinoma consistent with ovarian primary cells.
Applications:	This cell line is a suitable transfection host.
Tumor Formation::	Yes; Yes, in nude mice; forms moderately well differentiated adenocarcinoma consistent with ovarian primary
Products:	Antigen expression: Blood Type B; Rh+ Genes expressed: blood type B; Rh+ Isoenzymes: AK-1, 1 ES-D, 1 G6PD, B GLO-I, 1-2 Me-2, 1 PGM1, 1-2 PGM3, 1
Biosafety:	1

Data Sheet

PAGE 2/3

Additional Info:

SK-OV-3 cells are resistant to tumor necrosis factor and to several cytotoxic drugs including diphtheria toxin, cis-platinum and adriamycin.

Culture Medium:

McCoy's 5a + 2mM Glutamine + 15% of Fetal Bovine Serum (FBS).

Subculturing:

Remove and discard culture medium. Briefly rinse the cell layer with PBS without calcium and magnesium to remove all traces of serum which contains trypsin inhibitor. Add 2.0 to 3.0 mL of Trypsin-EDTA solution to flask and observe cells under an inverted microscope until cell layer is dispersed (usually within 5 to 15 minutes). Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal. Add 6.0 to 8.0 mL of complete growth medium and aspirate cells by gently pipetting. Add appropriate aliquots of the cell suspension to new culture vessels. Incubate cultures at 37°C.

**Subculturing Medium
Renewal:**

2 to 3 times per week

**Subculturing Subcultivation
Ratio:**

1:2 to 1:6

Culture Conditions:

Atmosphere: air, 95%; carbon dioxide (CO₂), 5% Temperature: 37°C

Cryopreservation:

95% FBS + 5% DMSO (Dimethyl sulfoxide)

SAFETY PRECAUTION: It is strongly recommended to always wear protective gloves, clothing, and a full-face mask when handling frozen vials. Some vials may leak when submerged in liquid nitrogen, allowing nitrogen to slowly enter the vial. Upon thawing, the conversion of liquid nitrogen back to its gas phase may cause the vial to explode or eject its cap with significant force, creating flying debris.

1. Thaw the vial by gently agitating it in a 37°C water bath. To minimize contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 minutes).
2. Remove the vial from the water bath as soon as its contents are thawed and decontaminate it by dipping in or spraying with 70% ethanol. From this point, all operations must be performed under strict aseptic conditions.
3. For cells sensitive to DMSO, it is recommended to remove the cryoprotective agent immediately. Transfer the vial contents to a centrifuge tube containing 9.0 mL of complete culture medium and centrifuge at approximately 125 × g for 5 to 7 minutes.
4. Discard the supernatant and resuspend the cell pellet in the recommended complete medium (see specific batch information for the appropriate dilution ratio).
5. Incubate the culture under appropriate atmospheric and temperature conditions (see "Culture Conditions" for this cell line).

NOTE: It is important to avoid excessive alkalinity of the medium during cell recovery. To minimize this risk, it is recommended to place the culture vessel containing the growth medium in the incubator for at least 15 minutes before adding the vial contents. This allows the medium to stabilize at its normal pH (7.0 to 7.6).

In Vitro 1975:155-159; J Nat Cancer Inst 1977;58:209-214 Abaan OD, Polley EC, Davis SR, Zhu YJ, Bilke S, Walker RL, Pineda M, Gindin Y, Jiang Y, Reinhold WC, Holbeck SL, Simon RM, Doroshow JH, Pommier Y, Meltzer PS. 2013 The exomes of the NCI-60 panel: a genomic resource for cancer biology and systems pharmacology. Cancer Res. 73(14):4372-82. PMID: 23856246.

Thawing Frozen Cells:

References:

Depositors:

Banco de Células do Rio de Janeiro

Cellosaurus:

[CVCL_0532](#)