

Data Sheet

PAGE 1/4

BCRJ Code:	0323
Cell Line:	T-47D
Species:	Homo sapiens
Vulgar Name:	Human
Tissue:	Mammary Gland; Derived From Metastatic Site: Pleural Effusion
Cell Type:	Epithelial
Morphology:	Epithelial
Disease:	Ductal Carcinoma
Growth Properties:	Adherent
Sex:	Female
Age/Ethnicity:	54 Year /
Derivation:	The T-47 line was isolated by I. Keydar from a pleural effusion obtained from a 54 year old female patient with an infiltrating ductal carcinoma of the breast.
Applications:	This cell line is a suitable transfection host.
Biosafety:	1
Additional Info:	This differentiated epithelial substrain (T-47D) was found to contain cytoplasmic junctions and receptors to 17 beta estradiol, other steroids and calcitonin. The cells express the WNT7B oncogene.
Culture Medium:	RPMI-1640 medium modified to contain 2 mM L-glutamine, 1 mM sodium pyruvate, 4500 mg/L glucose and 10% of fetal bovine serum.

Subculturing:

Volumes are given for a 75 cm² flask. Increase or decrease the amount of dissociation medium needed proportionally for culture vessels of other sizes. Corning® T-75 flasks (catalog #430641) are recommended for subculturing this product. Remove and discard culture medium. Briefly rinse the cell layer with PBS without calcium and magnesium to remove all traces of serum that contains trypsin inhibitor. Add 2.0 to 3.0 mL of Trypsin-EDTA solution to flask and observe cells under an inverted microscope until cell layer is dispersed (usually within 5 to 15 minutes). Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal. Add 6.0 to 8.0 mL of complete growth medium and aspirate cells by gently pipetting. Add appropriate aliquots of the cell suspension to new culture vessels. Incubate cultures at 37°C. Population Doubling Time 32 hrs NOTE: For more information on enzymatic dissociation and subculturing of cell lines consult Chapter 12 in Culture of Animal Cells, a manual of Basic Technique by R. Ian Freshney, 6th edition, published by Alan R. Liss, N.Y., 2010.

**Subculturing
Medium Renewal:**

2 to 3 times per week

**Subculturing
Subcultivation Ratio:**

1:3 to 1:5 is recommended

Culture Conditions:

Atmosphere: air, 95%; carbon dioxide (CO₂), 5% Temperature: 37°C

Cryopreservation:

95% FBS + 5% DMSO (Dimethyl sulfoxide)

Thawing Frozen Cells:

SAFETY PRECAUTION: It is highly recommended that protective gloves and clothing always be used and a full face mask always be worn when handling frozen vials. It is important to note that some vials leak when submerged in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vessel exploding or blowing off its cap with dangerous force creating flying debris.

1. Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the vial and cap out of the water. Thawing should be rapid (approximately 2 minutes).
2. Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
3. For cells that are sensitive to DMSO it is recommended that the cryoprotective agent be removed immediately. Transfer the vial contents to a centrifuge tube containing 9.0 mL complete culture medium and spin at approximately 125 x g for 5 to 7 minutes.
4. Discard the supernatant and Resuspend cell pellet with the recommended complete medium (see the specific batch information for the culture recommended dilution ratio).
5. Incubate the culture in an appropriate atmosphere and temperature (see "Culture Conditions" for this cell line).

NOTE: It is important to avoid excessive alkalinity of the medium during recovery of the cells. It is suggested that, prior to the addition of the vial contents, the culture vessel containing the growth medium be placed into the incubator for at least 15 minutes to allow the medium to reach its normal pH (7.0 to 7.6).

References:

Judge SM, Chatterton RT Jr.. Progesterone-specific stimulation of triglyceride biosynthesis in a breast cancer cell line (T-47D). *Cancer Res.* 43: 4407-4412, 1983. PubMed: 6871874 Lamp SJ, et al. Calcitonin induction of a persistent activated state of adenylate cyclase in human breast cancer cells (T-47D). *J. Biol. Chem.* 256: 12269-12274, 1981. PubMed: 6271778 Sher E, et al. Whole-cell uptake and nuclear localization of 1,25-dihydroxy-cholecalciferol by breast cancer cells (T-47D) in culture. *Biochem. J.* 200: 315-320, 1981. PubMed: 6896147 Freake HC, et al. 1,25-Dihydroxyvitamin D3 specifically binds to a human breast cancer cell line (T-47D) and stimulates growth. *Biochem. Biophys. Res. Commun.* 101: 1131-1138, 1981. PubMed: 6272774 Faust JB, Meeker TC. Amplification and expression of the bcl-1 gene in human solid tumor cell lines. *Cancer Res.* 52: 2460-2463, 1992. PubMed: 1568216 Huguet EL, et al. Differential expression of human Wnt genes 2, 3, 4, and 7B in human breast cell lines and normal and disease states of human breast tissue. *Cancer Res.* 54: 2615-2621, 1994. PubMed: 8168088 Bellet D, et al. Malignant transformation of nontrophoblastic cells is associated with the expression of chorionic gonadotropin beta genes normally transcribed in trophoblastic cells. *Cancer Res.* 57: 516-523, 1997. PubMed: 9012484 Hoppe HC, et al. Identification of phosphatidylinositol mannoside as a mycobacterial adhesin mediating both direct and opsonic binding to nonphagocytic mammalian cells. *Infect. Immun.* 65: 3896-3905, 1997. PubMed: 9284169 Burfeind P, et al. Antisense RNA to the type I insulin-like growth factor receptor suppresses tumor growth and prevents invasion by rat prostate cancer cells in vivo. *Proc. Natl. Acad. Sci. USA* 93: 7263-7268, 1996. PubMed: 8692980 Keydar I, et al. Establishment and characterization of a cell line of human breast carcinoma origin. *Eur. J. Cancer* 15: 659-670, 1979. PubMed: 228940

Depositors:

MARIA ISABEL ROSSI - UNIVERSIDADE FEDERAL DO RIO DE JANEIRO

ATCC:

HTB-133