

Data Sheet

PAGE 1/3

BCRJ Code:	0239
Cell Line:	U-138 MG
Species:	Homo sapiens
Vulgar Name:	Human
Tissue:	Brain
Morphology:	Polygonal
Disease:	Glioblastoma
Growth Properties:	Adherent
Sex:	Male
Age/Ethnicity:	47 Year / Caucasian
DNA Profile:	Amelogenin: X,Y CSF1PO: 12 D13S317: 9,11 D16S539: 12,13 D5S818: 11 D7S820: 9 TH01: 6 TPOX: 8 vWA: 18
Tumor Formation::	NO
Biosafety:	1
Culture Medium:	Dulbecco's Modified Eagle's Medium with 2 mM L-glutamine and fetal bovine serum to a final concentration of 10%.
Subculturing:	Remove medium, and rinse with PBS without calcium and magnesium. Remove the solution and add an additional 1 to 2 mL of trypsin-EDTA solution. Allow the flask to sit at room temperature (or at 37°C) until the cells detach. Add fresh culture medium, aspirate and dispense into new culture flasks. NOTE: For more information on enzymatic dissociation and subculturing of cell lines consult Chapter 12 in Culture of Animal Cells, a manual of Basic Technique by R. Ian Freshney, 6th edition, published by Alan R. Liss, N.Y., 2010.

Data Sheet

PAGE 2/3

Subculturing Medium Renewal:

2 to 3 times per week

Subculturing Subcultivation Ratio:

1:4 to 1:8

Culture Conditions:

Atmosphere: air, 95%; carbon dioxide (CO₂), 5% Temperature: 37°C

Cryopreservation:

95% FBS + 5% DMSO (Dimethyl sulfoxide)

Thawing Frozen Cells:

SAFETY PRECAUTION: It is strongly recommended to always wear protective gloves, clothing, and a full-face mask when handling frozen vials. Some vials may leak when submerged in liquid nitrogen, allowing nitrogen to slowly enter the vial. Upon thawing, the conversion of liquid nitrogen back to its gas phase may cause the vial to explode or eject its cap with significant force, creating flying debris.

1. Thaw the vial by gently agitating it in a 37°C water bath. To minimize contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 minutes).

2. Remove the vial from the water bath as soon as its contents are thawed and decontaminate it by dipping in or spraying with 70% ethanol. From this point, all operations must be performed under strict aseptic conditions.

3. For cells sensitive to DMSO, it is recommended to remove the cryoprotective agent immediately. Transfer the vial contents to a centrifuge tube containing 9.0 mL of complete culture medium and centrifuge at approximately 125 × g for 5 to 7 minutes.

4. Discard the supernatant and resuspend the cell pellet in the recommended complete medium (see specific batch information for the appropriate dilution ratio).

5. Incubate the culture under appropriate atmospheric and temperature conditions (see "Culture Conditions" for this cell line).

NOTE: It is important to avoid excessive alkalinity of the medium during cell recovery. To minimize this risk, it is recommended to place the culture vessel containing the growth medium in the incubator for at least 15 minutes before adding the vial contents. This allows the medium to stabilize at its normal pH (7.0 to 7.6).

Data Sheet

PAGE 3/3

References:

Beckman G , et al. G-6-PD and PGM phenotypes of 16 continuous human tumor cell lines. Evidence against cross-contamination and contamination by HeLa cells. Hum. Hered. 21: 238-241, 1971. Fogh J , et al. Absence of HeLa cell contamination in 169 cell lines

Depositors:

Ana Maria Oliveira Battastini, Universidade Federal Do Rio Grande Do Sul.

Cellosaurus:

[CVCL_0020](#)