

Banco de Células do Rio de Janeiro

Data Sheet

PAGE 1/3

BCRJ Code:	0240
Cell Line:	U266B1 [U266]
Species:	Homo sapiens
Vulgar Name:	Human
Tissue:	Peripheral Blood
Cell Type:	B Lymphocyte
Morphology:	Lymphoblast
Disease:	Myeloma; Plasmocytoma
Growth Properties:	Suspension
Age/Ethinicity:	53 Year /
Applications:	This cell line is a suitable transfection host.
DNA Profile:	Amelogenin: X,Y CSF1PO: 12,13 D13S317: 12 D16S539: 10 D5S818: 11,12 D7S820: 11,12 THO1: 5,7 TPOX: 8 vWA: 17
Products:	immunoglobulin; monoclonal antibody; interleukin 6 (interleukin-6, IL-6)
Biosafety:	1
Addtional Info:	U266 cells have been reported to produce human IL-6.
Culture Medium:	RPMI-1640 medium modified to contain 2 mM L-glutamine, 10 mM HEPES, 1 mM sodium pyruvate, 4500 mg/L glucose, and 1500 mg/L sodium bicarbonate and 15% of fetal bovine serum.

f

0



Banco de Células do Rio de Janeiro

Data	Sheet
------	-------

PAGE 2/3

Subculturing:	Cultures can be maintained by addition or replacement of fresh medium. Start cultures at 3 x 10e5 cells/mL and maintain between 1 x 10e5 and 1 x 10e6 cells/mL. T-75 flasks are recommended for subculturing this cell line.
Subculturing Medium Renewal:	2 to 3 times per week
Culture Conditions:	Atmosphere: air, 95%; carbon dioxide (CO2), 5% Temperature: 37°C
Cryopreservation:	95% FBS + 5% DMSO (Dimethyl sulfoxide)
Thawing Frozen Cells:	SAFETY PRECAUTION: It is strongly recommended to always wear protective gloves, clothing, and a full-face mask when handling frozen vials. Some vials may leak when submerged in liquid nitrogen, allowing nitrogen to slowly enter the vial. Upon thawing, the conversion of liquid nitrogen back to its gas phase may cause the vial to explode or eject its cap with significant force, creating flying debris. 1. Thaw the vial by gently agitating it in a 37°C water bath. To minimize contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 minutes). 2. Remove the vial from the water bath as soon as its contents are thawed and decontaminate it by dipping in or spraying with 70% ethanol. From this point, all operations must be performed under strict aseptic conditions. 3. For cells sensitive to DMSO, it is recommended to remove the cryoprotective agent immediately. Transfer the vial contents to a centrifuge tube containing 9.0 mL of complete culture medium and centrifuge at approximately $125 \times g$ for 5 to 7 minutes. 4. Discard the supernatant and resuspend the cell pellet in the recommended complete medium (see specific batch information for the appropriate dilution ratio). 5. Incubate the culture under appropriate atmospheric and temperature conditions (see "Culture Conditions" for this cell line). NOTE: It is important to avoid excessive alkalinity of the medium during cell recovery. To minimize this risk, it is recommended to place the culture vessel containing the growth medium in the incubator for at least 15 minutes before adding the vial contents. This allows the medium to stabilize at its normal pH (7.0 to 7.6).

f

0





Banco de Células do Rio de Janeiro

PAGE 3/3

References:	1180: Nilsson K, et al. Established immunoglobulin producing myeloma (IgE) and lymphoblastoid (IgG) cell lines from an IgE myeloma patient. Clin. Exp. Immunol. 7: 477-489, 1970. PubMed: 4097745 22377: Kawano M, et al. Autocrine generation and requirement
Depositors:	PRISCILA SEGGES; INCA
Cellosaurus:	<u>CVCL 0566</u>

f

