

Data Sheet

PAGE 1/3

BCRJ Code:	0243
Cell Line:	UMR-106
Species:	<i>Rattus norvegicus</i>
Vulgar Name:	Rat
Tissue:	Bone
Morphology:	Epithelial
Disease:	Osteosarcoma
Growth Properties:	Adherent
Derivation:	Both the original sarcoma and the cloned line were developed by T.J. Martin at the University of Sheffield. The UMR-106 cell line is a clonal derivative of a transplantable rat osteosarcoma that had been induced by injection of radiophosphorous (32P).
Biosafety:	1
Additional Info:	The cells are responsive to PTH, prostaglandins and bone resorbing steroids. Activation of protein kinase C inhibits ATP induced increases in intracellular calcium levels.
Culture Medium:	Dulbecco's Modified Eagle's Medium (DMEM) modified to contain 2 mM L-glutamine, 4500 mg/L glucose, 1 mM sodium pyruvate and 10% of fetal bovine serum.
Subculturing:	Remove medium, and rinse with 0.25% trypsin, 0.03% EDTA solution. Remove the solution and add an additional 1 to 2 mL of trypsin-EDTA solution. Allow the flask to sit at room temperature (or at 37°C) until the cells detach. Add fresh culture medium, aspirate and dispense into new culture flasks. NOTE: For more information on enzymatic dissociation and subculturing of cell lines consult Chapter 12 in Culture of Animal Cells, a manual of Basic Technique by R. Ian Freshney, 6th edition, published by Alan R. Liss, N.Y., 2010.

Data Sheet

PAGE 2/3

Subculturing Medium Renewal:

2 to 3 times per week

Subculturing Subcultivation Ratio:

1:4 to 1:8

Culture Conditions:

Atmosphere: air, 95%; carbon dioxide (CO₂), 5% Temperature: 37°C

Cryopreservation:

95% FBS + 5% DMSO (Dimethyl sulfoxide)

Thawing Frozen Cells:

SAFETY PRECAUTION: It is strongly recommended to always wear protective gloves, clothing, and a full-face mask when handling frozen vials. Some vials may leak when submerged in liquid nitrogen, allowing nitrogen to slowly enter the vial. Upon thawing, the conversion of liquid nitrogen back to its gas phase may cause the vial to explode or eject its cap with significant force, creating flying debris.

1. Thaw the vial by gently agitating it in a 37°C water bath. To minimize contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 minutes).
2. Remove the vial from the water bath as soon as its contents are thawed and decontaminate it by dipping in or spraying with 70% ethanol. From this point, all operations must be performed under strict aseptic conditions.
3. For cells sensitive to DMSO, it is recommended to remove the cryoprotective agent immediately. Transfer the vial contents to a centrifuge tube containing 9.0 mL of complete culture medium and centrifuge at approximately 125 × g for 5 to 7 minutes.
4. Discard the supernatant and resuspend the cell pellet in the recommended complete medium (see specific batch information for the appropriate dilution ratio).
5. Incubate the culture under appropriate atmospheric and temperature conditions (see "Culture Conditions" for this cell line).

NOTE: It is important to avoid excessive alkalinity of the medium during cell recovery. To minimize this risk, it is recommended to place the culture vessel containing the growth medium in the incubator for at least 15 minutes before adding the vial contents. This allows the medium to stabilize at its normal pH (7.0 to 7.6).

References:

Nature (Lond.) 260: 436-438, 1976; Eur. J. Cancer 15: 1151-1158, 1979; Clin. Orthop. Rel. Res. 140: 247-254, 1979; FEBS Lett. 115: 139-142, 1980; Cancer Res. 43: 4308-4315, 1983; Methods Enzymol. 145: 324-336, 1987;

Data Sheet

PAGE 3/3

Depositors:

Dr. Willian George Goodman, Department of Radiology, UCLA School of Medicine through Dr. Maria Eugênia Leite Duarte, Universidade Federal Fluminense.

Cellosaurus:

[CVCL 3617](#)